

# PASSIVE SYSTEMS WITH A NORMAL MAIN OPERATOR AND QUASI-SELFADJOINT SYSTEMS

YU.M. ARLINSKIĬ, S. HASSI, AND H.S.V. DE SNOO

**ABSTRACT.** Passive systems  $\tau = \{T, \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  with  $\mathfrak{M}$  and  $\mathfrak{N}$  as an input and output space and  $\mathfrak{H}$  as a state space are considered in the case that the main operator on the state space is normal. Basic properties are given and a general unitary similarity result involving some spectral theoretic conditions on the main operator is established. A passive system  $\tau$  with  $\mathfrak{M} = \mathfrak{N}$  is said to be quasi-selfadjoint if  $\text{ran}(T - T^*) \subset \mathfrak{N}$ . The subclass  $\mathbf{S}^{qs}(\mathfrak{N})$  of the Schur class  $\mathbf{S}(\mathfrak{N})$  is the class formed by all transfer functions of quasi-selfadjoint passive systems. The subclass  $\mathbf{S}^{qs}(\mathfrak{N})$  is characterized and minimal passive quasi-selfadjoint realizations are studied. The connection between the transfer function belonging to the subclass  $\mathbf{S}^{qs}(\mathfrak{N})$  and the  $Q$ -function of  $T$  is given.

## 1. INTRODUCTION

Let  $\mathfrak{M}$ ,  $\mathfrak{N}$ , and  $\mathfrak{H}$  be separable Hilbert spaces and let

$$(1.1) \quad T = \begin{pmatrix} D & C \\ B & A \end{pmatrix} : \begin{pmatrix} \mathfrak{M} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix}$$

be a bounded linear operator. Here and in the following, it will be tacitly assumed that the spaces in the righthand side are orthogonal sums:  $\mathfrak{M} \oplus \mathfrak{H}$  and  $\mathfrak{N} \oplus \mathfrak{H}$ . The system of equations

$$(1.2) \quad \begin{cases} h_{k+1} = Ah_k + B\xi_k, \\ \sigma_k = Ch_k + D\xi_k, \end{cases} \quad k \geq 0,$$

describes the evolution of a *linear discrete time-invariant system*  $\tau = \{T, \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$ . The Hilbert spaces  $\mathfrak{M}$  and  $\mathfrak{N}$  are called the input and the output spaces, respectively, and the Hilbert space  $\mathfrak{H}$  is called the state space. The operators  $A$ ,  $B$ ,  $C$ , and  $D$  are called the *main operator*, the *control operator*, the *observation operator*, and the *feedthrough operator* of  $\tau$ , respectively. The subspaces

$$(1.3) \quad \mathfrak{H}^c = \overline{\text{span}} \{ A^n B \mathfrak{M} : n \in \mathbb{N}_0 \} \text{ and } \mathfrak{H}^o = \overline{\text{span}} \{ A^{*n} C^* \mathfrak{N} : n \in \mathbb{N}_0 \}$$

are called the *controllable* and *observable* subspaces of  $\tau = \{T, \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$ , respectively. If  $\mathfrak{H}^c = \mathfrak{H}$  ( $\mathfrak{H}^o = \mathfrak{H}$ ) then the system  $\tau$  is said to be *controllable* (*observable*), and *minimal* if  $\tau$  is both controllable and observable. If  $\mathfrak{H} = \text{clos} \{ \mathfrak{H}^c + \mathfrak{H}^o \}$  then the system  $\tau$  is said to be a *simple*. Two discrete-time systems  $\tau_1 = \{T_1, \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_1\}$  and

---

*Key words and phrases.* Passive system, transfer function, quasi-selfadjoint contraction,  $Q$ -function.

This work was partially supported by the Research Institute for Technology at the University of Vaasa. The first author was also supported by the Academy of Finland (projects 117617, 123902) and the Dutch Organization for Scientific Research NWO (B 61-553).

$\tau_2 = \{T_2, \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_2\}$  are *unitarily similar* if there exists a unitary operator  $U$  from  $\mathfrak{H}_1$  onto  $\mathfrak{H}_2$  such that

$$(1.4) \quad A_2 = UA_1U^*, \quad B_2 = UB_1, \quad C_2 = C_1U^*, \quad \text{and} \quad D_2 = D_1.$$

If the linear operator  $T$  is contractive (isometric, co-isometric, unitary), then the corresponding discrete-time system is said to be *passive* (*isometric*, *co-isometric*, *conservative*). The *transfer function*

$$(1.5) \quad \Theta(\lambda) := D + \lambda C(I - \lambda A)^{-1}B, \quad \lambda \in \mathbb{D},$$

of the passive system  $\tau$  in (1.2) belongs to the *Schur class*  $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ , i.e.,  $\Theta(\lambda)$  is holomorphic in the unit disk  $\mathbb{D} = \{\lambda \in \mathbb{C} : |\lambda| < 1\}$  and its values are contractive linear operators from  $\mathfrak{M}$  into  $\mathfrak{N}$ . Every operator-valued function  $\Theta(\lambda)$  from the Schur class  $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$  can be realized as the transfer function of a passive system, which can be chosen as observable co-isometric (controllable isometric, simple conservative, passive minimal). Moreover two isometric and controllable (co-isometric and observable, simple conservative) systems having the same transfer function are unitarily similar. D.Z. Arov [10] has shown that two minimal passive systems  $\tau_1$  and  $\tau_2$  with the same transfer function  $\Theta(\lambda)$  are only *weakly similar*, i.e., there is a closed densely defined operator  $Z : \mathfrak{H}_1 \rightarrow \mathfrak{H}_2$  such that  $Z$  is invertible,  $Z^{-1}$  is densely defined, and

$$(1.6) \quad ZA_1f = A_2Zf, \quad C_1f = C_2Zf, \quad f \in \text{dom } Z, \quad \text{and} \quad ZB_1 = B_2.$$

Weak similarity preserves neither the dynamical properties of the system nor the spectral properties of its main operator  $A$ . In [14], [15] necessary and sufficient conditions have been established for minimal passive systems with the same transfer function to be (unitarily) similar. In [5] a parametrization of the contractive block-operator matrices in (1.1) was used to establish some new aspects and some explicit formulas for the interplay between the system  $\tau$ , its transfer function  $\Theta(\lambda)$ , and the Sz.-Nagy–Foiaş characteristic function of the contraction  $A$ .

In this paper the same approach is applied to study passive systems with a normal main operator, including the class of *passive quasi-selfadjoint* systems (*pqs*-systems for short), as defined in the paper. Furthermore, using the famous Mergelyan's theorem from complex analysis a general unitary similarity result is proved for such systems.

The passive system  $\tau = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  is called a *pqs*-system if the operator

$$(1.7) \quad T = \begin{pmatrix} D & C \\ B & A \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix}$$

is a *quasi-selfadjoint contraction* (*qsc*-operator for short), i.e.,  $T$  is a contraction and  $\text{ran}(T - T^*) \subset \mathfrak{N}$ , cf. [4]. This last condition is equivalent to  $A = A^*$  and  $C = B^*$ . If  $\tau$  is a *pqs*-system, then the transfer function (1.5) of  $\tau$  takes the form

$$\Theta(\lambda) = W(\lambda) + D,$$

where the function  $W(\lambda)$  is a Herglotz–Nevanlinna function defined on  $\text{Ext}\{(-\infty, -1] \cup [1, \infty)\}$ . The subclass  $\mathbf{S}^{qs}(\mathfrak{N})$  of the Schur class  $\mathbf{S}(\mathfrak{N})$  of  $\mathbf{L}(\mathfrak{N})$ -valued functions is the class of all transfer functions of *pqs*-systems  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$ . A necessary and sufficient condition for the function  $\Theta(\lambda)$  to be in the class  $\mathbf{S}^{qs}$  is given, the minimal *pqs*-systems with the given operator-valued function  $\Theta(\lambda)$  from the class  $\mathbf{S}^{qs}$  is constructed using operator representations of Herglotz–Nevanlinna functions. Moreover, a necessary

and sufficient condition for the function  $\Theta(\lambda) \in \mathbf{S}^{qs}$  to be inner (co-inner) is proved and connections with  $pqs$ -system and other minimal systems with the same transfer function are established. Also it is shown that if, for instance,  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  and  $\varphi_\Theta(\lambda) = 0$  ( $\psi_\Theta(\lambda) = 0$ ) then  $\Theta(\lambda)$  is inner (co-inner). A matrix form of the inner function from the class  $\mathbf{S}^{qs}(\mathfrak{N})$  when  $\dim \mathfrak{N} < \infty$  is also given, and in the case of scalar functions from the class  $\mathbf{S}^{qs}(\mathfrak{N})$  a minimal representation is obtained by means of Jacobi matrices.

## 2. PRELIMINARIES

Let  $\mathfrak{M}$  and  $\mathfrak{N}$  be Hilbert spaces and let  $\Theta(\lambda)$  belong to the Schur class  $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ . The notation  $\Theta(\xi)$ ,  $\xi \in \mathbb{T}$ , stands for the non-tangential strong limit value of  $\Theta(\lambda)$  which exist almost everywhere on  $\mathbb{T}$ , cf. [27]. A function  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  is said to be *inner* if  $\Theta^*(\xi)\Theta(\xi) = I_{\mathfrak{M}}$  for almost all  $\xi \in \mathbb{T}$ , and it is said to be *co-inner* if  $\Theta(\xi)\Theta^*(\xi) = I_{\mathfrak{N}}$  for almost all  $\xi \in \mathbb{T}$ . A function  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  is said to be *bi-inner* if it is both inner and co-inner.

**2.1. Contractions and their defect operators.** Let  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  be a contraction, in other words, let  $\|Af\| \leq \|f\|$  for all  $f \in \mathfrak{H}$ , or equivalently  $I - A^*A \geq 0$ . The selfadjoint operator  $D_A = (I - A^*A)^{1/2}$  is said to be the *defect operator* of  $A$ . Observe that  $\ker D_A = \ker D_A^2 = \ker (I - A^*A)$ , and that

$$(2.1) \quad \ker (I - A^*A) = \{ f \in \mathfrak{H} : \|Af\| = \|f\| \}.$$

Clearly, any contraction  $A$  satisfies

$$I \geq A^*A \geq \cdots \geq A^{*n}A^n \geq 0, \quad n \in \mathbb{N},$$

in other words the sequence  $\|A^n f\|$  with  $f \in \mathfrak{H}$  is monotonically nonincreasing. In particular, the strong limit

$$(2.2) \quad S_A = s - \lim A^{*n}A^n,$$

exists as an operator in  $\mathbf{L}(\mathfrak{H}_1)$ , cf. [25, p. 261]. The defect operators  $D_A$  and  $D_{A^*}$  satisfy the following commutation relation:

$$(2.3) \quad AD_A = D_{A^*}A, \quad D_AA^* = A^*D_{A^*}.$$

Let  $\mathfrak{D}_A$  stand for the closure of the range  $\text{ran } D_A$ . Then

$$(2.4) \quad \begin{pmatrix} -A & D_{A^*} \\ D_A & A^* \end{pmatrix} : \begin{pmatrix} \mathfrak{D}_A \\ \mathfrak{H}_2 \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{D}_{A^*} \\ \mathfrak{H}_1 \end{pmatrix}$$

is unitary. Define the subspaces  $\mathfrak{H}_{A,0}$  and  $\mathfrak{H}_{A,1}$  by

$$\mathfrak{H}_{A,1} = \{ f \in \mathfrak{H} : \|f\| = \|A^n f\| = \|A^{*n} f\|, \quad n \in \mathbb{N} \}, \quad \mathfrak{H}_{A,0} = \mathfrak{H} \ominus \mathfrak{H}_{A,1}.$$

Then  $\mathfrak{H} = \mathfrak{H}_{A,0} \oplus \mathfrak{H}_{A,1}$  is a canonical orthogonal decomposition of  $\mathfrak{H}$  such that

$$(2.5) \quad A = A_0 \oplus A_1, \quad A_j = A|_{\mathfrak{H}_{A,j}}, \quad j = 0, 1,$$

where  $\mathfrak{H}_{A,0}$  and  $\mathfrak{H}_{A,1}$  reduce  $A$ ,  $A_0$  is a completely non-unitary contraction, and  $A_1$  is a unitary operator. The function

$$(2.6) \quad \Phi_A(\lambda) := (-A + \lambda D_{A^*}(I_{\mathfrak{H}} - \lambda A^*)^{-1}D_A)|_{\mathfrak{D}_A}, \quad \lambda \in \mathbb{D},$$

is the Sz.-Nagy–Foiaş characteristic function of the contraction  $A$ . It belongs to the Schur class  $\mathbf{S}(\mathfrak{D}_A, \mathfrak{D}_{A^*})$ ; cf. [27]. In fact, a straightforward calculation using the identities  $(I - \lambda A)^{-1} = I + \lambda(I - \lambda A)^{-1}A$  and (2.3) yields

$$(2.7) \quad D_{\Phi_A(\lambda)}^2 = (1 - \lambda \bar{\lambda}) D_A (I - \bar{\lambda} A)^{-1} (I - \lambda A^*)^{-1} D_A \upharpoonright \mathfrak{D}_A,$$

which shows that  $\Phi_A(\lambda)$  is contractive for  $\lambda \in \mathbb{D}$ . Note also that  $\Phi_{A^*}(\lambda)$  is the transfer function of the conservative system

$$(2.8) \quad \Sigma = \{T, \mathfrak{D}_{A^*}, \mathfrak{D}_A, \mathfrak{H}\},$$

where

$$(2.9) \quad T = \begin{pmatrix} -A^* & D_A \\ D_{A^*} & A \end{pmatrix} : \begin{pmatrix} \mathfrak{D}_{A^*} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{D}_A \\ \mathfrak{H} \end{pmatrix}.$$

Let  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  be a contraction and let  $\Sigma$  be the corresponding conservative system in (2.8), (2.9). Then the controllable and observable subspaces, as defined in (1.3), are given by

$$(2.10) \quad \mathfrak{H}_\Sigma^c = \overline{\text{span}} \{A^n D_{A^*} \mathfrak{D}_{A^*} : n \in \mathbb{N}_0\}, \quad \mathfrak{H}_\Sigma^o = \overline{\text{span}} \{A^{*n} D_A \mathfrak{D}_A : n \in \mathbb{N}_0\}.$$

Observe that  $A$  is completely nonunitary if and only if  $\Sigma$  is minimal. Since clearly  $\Phi_A(\lambda)^* = \Phi_{A^*}(\bar{\lambda})$ , one has also

$$(2.11) \quad D_{\Phi_A(\lambda)^*}^2 = (1 - \bar{\lambda} \lambda) D_{A^*} (I - \lambda A^*)^{-1} (I - \bar{\lambda} A)^{-1} D_{A^*} \upharpoonright \mathfrak{D}_{A^*}.$$

Observe that if  $\xi \in \mathbb{T} := \{\xi \in \mathbb{C} : |\xi| = 1\}$  belongs to the resolvent set of  $A$ , then (2.7) and (2.11) show that  $\Phi_A(\xi)$  is a unitary operator; cf. [27, p. 239].

A contraction  $A$  in a Hilbert space  $\mathfrak{H}$  is said to belong to the *classes*  $C_0$ . or  $C_{0.}$  if

$$s - \lim_{n \rightarrow \infty} A^n = 0 \quad \text{or} \quad s - \lim_{n \rightarrow \infty} A^{*n} = 0,$$

respectively. By definition,  $C_{00} := C_0. \cap C_{0.}$ . Hence  $A \in C_{00}$  precisely when

$$(2.12) \quad s - \lim_{n \rightarrow \infty} A^n = s - \lim_{n \rightarrow \infty} A^{*n} = 0.$$

Observe that  $A \in C_{00}$  implies that  $A$  is completely nonunitary, cf. (2.5). The completely non-unitary part of a contraction  $A$  belongs to the class  $C_{0.}$ ,  $C_{0.}$ , or  $C_{00}$  if and only if its characteristic function  $\Phi_A(\lambda)$  in (2.6) is inner, co-inner, or bi-inner, respectively; cf. [27, Theorem VI.2.3]. It follows from (2.2) that  $S_A = 0$  implies  $A \in C_{0.}$  and that  $S_{A^*} = 0$  implies  $A \in C_{0.}$ .

A contraction  $A$  is said to be *strict* if  $\|Af\| < \|f\|$  for all nontrivial  $f \in \mathfrak{H}_1$ . Note that in view of (2.1) a contraction  $A$  is strict if and only if  $\ker D_A = \ker D_A^2 = \ker (I - A^*A) = \{0\}$ . Finally, a passive system  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  is said to be *strongly stable* or *strongly co-stable* if the main operator  $A$  belongs to the class  $C_{0.}$  or  $C_{0.}$ , respectively; see [11], [16].

**2.2. Some properties of normal contractions.** An operator  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  is said to be *normal* if  $A^*A = AA^*$ , or equivalently, if  $\|Af\| = \|A^*f\|$  for all  $f \in \mathfrak{H}$ , cf. [25, p. 281]. It is clear from  $\mathfrak{H} = \mathfrak{H}_{A,0} \oplus \mathfrak{H}_{A,1}$  and the orthogonal decomposition in (2.5) that

a contraction  $A$  is normal if and only if its completely nonunitary part  $A_0$  is normal in  $\mathfrak{H}_{A,0}$ . If  $A$  is a normal contraction then, parallel to (2.1), one has

$$(2.13) \quad \begin{aligned} \ker (I - (A^*A)^n) &= \ker (I - A^{*n}A^n) = \{ f \in \mathfrak{H} : \|A^n f\| = \|f\| \} \\ &= \ker (I - (AA^*)^n) = \ker (I - A^n A^{*n}) = \{ f \in \mathfrak{H} : \|A^{*n} f\| = \|f\| \}. \end{aligned}$$

Moreover, if  $A$  is a normal contraction, then the defect operators  $D_A$  and  $D_{A^*}$  satisfy  $D_A = D_{A^*}$  and  $\mathfrak{D}_A = \mathfrak{D}_{A^*}$ ; in addition, (2.3) reads as

$$(2.14) \quad AD_A = D_A A, \quad A^* D_A = D_A A^*.$$

**Lemma 2.1.** *Let  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  be a normal contraction. Then the strong limit  $S_A$  satisfies  $S_A = S_{A^*}$  and*

$$(2.15) \quad S_A(I - A^*A) = 0.$$

*If, in addition,  $A$  is strict, then  $S_A = 0$ .*

*Proof.* If  $A$  is normal, then (2.2) implies that  $S_A = S_{A^*}$  and

$$S_A A^* A = (s - \lim A^{*n} A^n) A^* A = s - \lim A^{*(n+1)} A^{n+1} = S_A,$$

which leads to (2.15).  $\square$

**Proposition 2.2.** *Let  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  be a normal contraction. Then the following statements are equivalent:*

- (i)  $A \in C_{00}$ ;
- (ii)  $A$  is completely non-unitary;
- (iii)  $A$  is strict.

*Moreover, the characteristic function  $\Phi_A(\lambda)$  of  $A$  in (2.6) is bi-inner.*

*Proof.* (i)  $\Rightarrow$  (ii) This implication is a general fact for not necessarily normal contractions.

(ii)  $\Rightarrow$  (iii) Let  $A$  be completely non-unitary. Assume that  $A$  is not strict. Then there exists an element  $0 \neq f_0 \in \mathfrak{H}_1$  such that  $\|A f_0\| = \|f_0\|$ . Since  $\ker (I - A^*A) \subset \ker (I - (A^*A)^n)$ ,  $n \in \mathbb{N}$ , it follows from (2.1) and (2.13) that  $\|f_0\| = \|A^n f_0\| = \|A^{*n} f_0\| > 0$ . This contradicts the fact that  $A$  is completely nonunitary.

(iii)  $\Rightarrow$  (i) Let  $A$  be strict, so that  $\ker (I - A^*A) = \{0\}$ . Then Lemma 2.1 implies that  $S_A = 0$ , which leads to (2.12), so that  $A \in C_{00}$ .

Observe that if  $A$  is normal then  $\mathfrak{H}_{A,1} = \ker D_A$  as was just shown above. The completely non-unitary part  $A_0$  of  $A$  is normal and satisfies  $\ker D_{A_0} = \{0\}$ . Thus  $A_0 \in C_{00}$ , i.e.,  $\Phi_A(\lambda)$  is bi-inner; cf. [27, Theorem VI.2.3].  $\square$

If a contraction  $A$  is normal, then its controllable and observable subspaces coincide, which leads to the following observation.

**Proposition 2.3.** *Let  $A \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$  be a normal contraction, and let  $\Sigma$  be the corresponding conservative system in (2.8), (2.9). Then  $\mathfrak{H}_\Sigma^c = \mathfrak{H}_\Sigma^o$  and the following statements are equivalent:*

- (i)  $\Sigma$  is simple;
- (ii)  $\Sigma$  is controllable;
- (iii)  $\Sigma$  is observable;
- (iv)  $\Sigma$  is minimal.

*Proof.* Since  $A$  is normal, it follows that  $D_{A^{*n}} = D_{A^n}$  for all  $n \in \mathbb{N}_0$ . Hence the identities

$$(2.16) \quad (\mathfrak{H}_\Sigma^c)^\perp = \bigcap_{n=0}^{\infty} \ker (D_{A^*} A^{*n}) = \bigcap_{n=1}^{\infty} \ker D_{A^{*n}},$$

$$(2.17) \quad (\mathfrak{H}_\Sigma^o)^\perp = \bigcap_{n=0}^{\infty} \ker (D_A A^n) = \bigcap_{n=1}^{\infty} \ker D_{A^n}$$

imply that  $(\mathfrak{H}_\Sigma^c)^\perp = (\mathfrak{H}_\Sigma^o)^\perp$ , or equivalently,  $\mathfrak{H}_\Sigma^c = \mathfrak{H}_\Sigma^o$ . This identity implies the equivalence of (i), (ii), (iii), and (iv).  $\square$

The following corollary is based on the fact that a contraction  $A$  is completely nonunitary if and only if the corresponding system  $\Sigma$  in (2.8), (2.9) is minimal.

**Corollary 2.4.** *Let  $A$  be a normal contraction. Then the statements (i)–(iii) in Proposition 2.2 and the statements (i)–(iv) in Proposition 2.3 are all equivalent.*

**2.3. Parametrization of block operators.** For a proof and some history of the following theorem, see [5].

**Theorem 2.5.** *Let  $\mathfrak{M}, \mathfrak{N}, \mathfrak{H}$ , and  $\mathfrak{K}$  be Hilbert spaces. The operator matrix  $T$  in (1.1) is a contraction if and only if  $T$  is of the form*

$$(2.18) \quad T = \begin{pmatrix} -KA^*M + D_{K^*}XD_M & KD_A \\ D_{A^*}M & A \end{pmatrix},$$

where  $A \in \mathbf{L}(\mathfrak{H}, \mathfrak{K})$ ,  $M \in \mathbf{L}(\mathfrak{M}, \mathfrak{D}_{A^*})$ ,  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$ , and  $X \in \mathbf{L}(\mathfrak{D}_M, \mathfrak{D}_{K^*})$  are contractions, all uniquely determined by  $T$ . Furthermore, the following equality holds for all  $h \in \mathfrak{M}, f \in \mathfrak{H}$ :

$$(2.19) \quad \left\| \begin{pmatrix} h \\ f \end{pmatrix} \right\|^2 - \left\| \begin{pmatrix} -KA^*M + D_{K^*}XD_M & KD_A \\ D_{A^*}M & A \end{pmatrix} \begin{pmatrix} h \\ f \end{pmatrix} \right\|^2 \\ = \|D_K(D_A f - A^* M h) - K^* X D_M h\|^2 + \|D_X D_M h\|^2.$$

**Corollary 2.6.** *Let  $A \in \mathbf{L}(\mathfrak{H}, \mathfrak{K})$  be a contraction. Assume that  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$ ,  $M \in \mathbf{L}(\mathfrak{M}, \mathfrak{D}_{A^*})$ , and  $X \in \mathbf{L}(\mathfrak{D}_M, \mathfrak{D}_{K^*})$  are contractions. Then the operator  $T$  in (2.18) is:*

- (i) *isometric if and only if  $D_X D_M = 0$  and  $D_K D_A = 0$ ;*
- (ii) *co-isometric if and only if  $D_{X^*} D_{K^*} = 0$  and  $D_{M^*} D_{A^*} = 0$ .*

Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system and let (2.18) be the representation of the block operator  $T$  in (1.1). Define for  $\lambda \in \mathbb{D}$  the following operator-valued holomorphic functions

$$(2.20) \quad \varphi(\lambda) := \begin{pmatrix} -D_X D_M \\ D_K \Phi_{A^*}(\lambda) M - K^* X D_M \end{pmatrix} : \mathfrak{M} \rightarrow \begin{pmatrix} \mathfrak{D}_M \\ \mathfrak{D}_K \end{pmatrix},$$

and

$$(2.21) \quad \psi(\lambda) := \begin{pmatrix} D_{K^*} D_{X^*} & K \Phi_{A^*}(\lambda) D_{M^*} - D_{K^*} X M^* \end{pmatrix} : \begin{pmatrix} \mathfrak{D}_{K^*} \\ \mathfrak{D}_{M^*} \end{pmatrix} \rightarrow \mathfrak{N}.$$

**Theorem 2.7** ([5]). *Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system and let (2.18) be the representation of the block operator  $T$  in (1.1). Then the transfer function  $\Theta(\lambda)$  of  $\tau$  and the characteristic function  $\Phi_{A^*}(\lambda)$  of  $A^*$  (see (2.6)) are connected via*

$$(2.22) \quad \Theta(\lambda) = K\Phi_{A^*}(\lambda)M + D_{K^*}XD_M, \quad \lambda \in \mathbb{D};$$

in particular,  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ . In addition, the identities

$$(2.23) \quad \|D_{\Theta(\lambda)}h\|^2 = \|D_{\Phi_{A^*}(\lambda)}Mh\|^2 + \|\varphi(\lambda)h\|^2, \quad h \in \mathfrak{M},$$

$$(2.24) \quad \|D_{\Theta^*(\lambda)}g\|^2 = \|D_{\Phi_A(\bar{\lambda})}K^*g\|^2 + \|\psi^*(\lambda)g\|^2, \quad g \in \mathfrak{N},$$

hold and the functions  $\varphi(\lambda)$  and  $\psi(\lambda)$  in (2.20) and (2.21) are Schur functions.

### 3. PASSIVE SYSTEMS WITH A NORMAL MAIN OPERATOR

Let  $\tau$  be a passive system of the form (1.1). If its main operator  $A$  is normal, then many properties of  $\tau$  and its transfer function simplify.

**3.1. Basic properties.** The controllable and observable subspaces of the passive system in (1.1) are defined in (1.3). Let the block matrix  $T$  have the parametrization (2.18), so that  $A^n B = A^n D_{A^*} M$  and  $A^{*n} C^* = A^{*n} D_A K^*$ . If, in addition,  $A$  is normal it follows that  $D_{A^*} = D_A$  and then (2.14) implies

$$A^n B = D_A A^n M, \quad A^{*n} C^* = A^{*n} D_A K^*.$$

Hence, if  $A$  is normal, then  $\mathfrak{H}^c$  and  $\mathfrak{H}^o$  have the form:

$$(3.1) \quad \mathfrak{H}^c = \overline{\text{span}} \{ D_A A^n M \mathfrak{M} : n \in \mathbb{N}_0 \}, \quad \mathfrak{H}^o = \overline{\text{span}} \{ D_A A^{*n} K^* \mathfrak{N} : n \in \mathbb{N}_0 \},$$

or, equivalently,

$$(3.2) \quad (\mathfrak{H}^c)^\perp = \bigcap_{n=0}^{\infty} \ker (M^* A^{*n} D_A), \quad (\mathfrak{H}^o)^\perp = \bigcap_{n=0}^{\infty} \ker (K A^n D_A),$$

Let the subspaces  $\mathfrak{H}_N^c$  and  $\mathfrak{H}_N^o$  be defined by

$$(3.3) \quad \mathfrak{H}_N^c = \overline{\text{span}} \{ A^n M \mathfrak{M} : n \in \mathbb{N}_0 \}, \quad \mathfrak{H}_N^o = \overline{\text{span}} \{ A^{*n} K^* \mathfrak{N} : n \in \mathbb{N}_0 \},$$

or, equivalently, by

$$(3.4) \quad (\mathfrak{H}_N^c)^\perp = \bigcap_{n=0}^{\infty} \ker (M^* A^{*n}), \quad (\mathfrak{H}_N^o)^\perp = \bigcap_{n=0}^{\infty} \ker (K A^n).$$

**Lemma 3.1.** *Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system with  $T$  of the form (2.18) with some contractions  $A \in \mathbf{L}(\mathfrak{H}, \mathfrak{H})$ ,  $M \in \mathbf{L}(\mathfrak{M}, \mathfrak{D}_{A^*})$ ,  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$ , and  $X \in \mathbf{L}(\mathfrak{D}_M, \mathfrak{D}_{K^*})$ . Assume that  $A$  is normal.*

- (i) *If  $\mathfrak{H}_N^c$  is invariant under  $A^*$ , then  $\mathfrak{H} \ominus \mathfrak{H}_N^c \subset \mathfrak{H} \ominus \mathfrak{H}^c$ .*
- (ii) *If  $\mathfrak{H}_N^o$  is invariant under  $A$ , then  $\mathfrak{H} \ominus \mathfrak{H}_N^o \subset \mathfrak{H} \ominus \mathfrak{H}^o$ .*

*Proof.* (i) Assume that  $\mathfrak{H}_N^c$  is invariant under  $A^*$  or, equivalently, that  $\mathfrak{H} \ominus \mathfrak{H}_N^c$  is invariant under  $A$ . Hence, if  $f \in \mathfrak{H} \ominus \mathfrak{H}_N^c$  then  $f$  and  $Af$  both belong to  $\ker (M^* A^{*n})$  for all  $n \in \mathbb{N}_0$ . Thus, in particular,  $D_A^2 f = (I - A^* A)f \in \ker M^*$ . Moreover, if  $p(t)$  is a polynomial then  $p(D_A^2)f \in \ker M^*$ . Since there exists a sequence of polynomials  $\{p_m(t)\}_{m=1}^\infty$  such that the sequence  $\{p_m(D_A^2)\}$  converges uniformly to  $D_A$ , it follows that  $D_A f \in \ker M^*$ . Furthermore, the sequence  $\{p_m(D_A^2)A^{*n}\}$  converges uniformly to  $D_A A^{*n}$  for all  $n \in \mathbb{N}$ . Since  $p_m(D_A^2)A^{*n}f \in \ker M^*$  for all  $n \in \mathbb{N}_0$ , one concludes that  $D_A A^{*n}f \in \ker M^*$  for all  $n \in \mathbb{N}_0$ . It follows that

$$\mathfrak{H} \ominus \mathfrak{H}_N^c \subset \bigcap_{n=0}^{\infty} \ker (M^* D_A A^{*n}) = \mathfrak{H} \ominus \overline{\text{span}} \{ D_A A^n M \mathfrak{N} : n \in \mathbb{N}_0 \} = \mathfrak{H} \ominus \mathfrak{H}^c.$$

(ii) The proof of (ii) is similar to the proof of (i).  $\square$

**Proposition 3.2.** *Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system where  $T$  is of the form (2.18) with contractions  $A \in \mathbf{L}(\mathfrak{H}, \mathfrak{K})$ ,  $M \in \mathbf{L}(\mathfrak{M}, \mathfrak{D}_{A^*})$ ,  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$ , and  $X \in \mathbf{L}(\mathfrak{D}_M, \mathfrak{D}_{K^*})$ . Assume that  $A$  is normal.*

(i)  $\tau$  is controllable if and only if

$$(3.5) \quad \ker D_A = \{0\} \quad \text{and} \quad \text{ran } D_A \cap (\mathfrak{H} \ominus \mathfrak{H}_N^c) = \{0\}.$$

*In particular, if  $\ker D_A = \{0\}$  and  $\mathfrak{H}_N^c = \mathfrak{H}$ , then  $\tau$  is controllable; if  $\tau$  is controllable and  $\mathfrak{H}_N^c$  is invariant under  $A^*$ , then  $\ker D_A = \{0\}$  and  $\mathfrak{H}_N^c = \mathfrak{H}$ .*

(ii)  $\tau$  is observable if and only if

$$(3.6) \quad \ker D_A = \{0\} \quad \text{and} \quad \text{ran } D_A \cap (\mathfrak{H} \ominus \mathfrak{H}_N^o) = \{0\};$$

*In particular, if  $\ker D_A = \{0\}$  and  $\mathfrak{H}_N^o = \mathfrak{H}$ , then  $\tau$  is observable; if  $\tau$  is observable and  $\mathfrak{H}_N^o$  is invariant under  $A$ , then  $\ker D_A = \{0\}$  and  $\mathfrak{H}_N^o = \mathfrak{H}$ .*

(iii)  $\tau$  is simple if and only if

$$(3.7) \quad \ker D_A = \{0\} \quad \text{and} \quad \text{ran } D_A \cap \mathfrak{H} \ominus (\mathfrak{H}_N^c + \mathfrak{H}_N^o) = \{0\}.$$

*In particular, if  $\ker D_A = \{0\}$  and  $\mathfrak{H} = \text{clos} \{\mathfrak{H}_N^c + \mathfrak{H}_N^o\}$ , then  $\tau$  is simple; if  $\tau$  is simple,  $\mathfrak{H}_N^c$  is invariant under  $A^*$ , and  $\mathfrak{H}_N^o$  is invariant under  $A$ , then  $\ker D_A = \{0\}$  and  $\mathfrak{H} = \text{clos} \{\mathfrak{H}_N^c + \mathfrak{H}_N^o\}$ .*

(iv)  $\tau$  is minimal if and only if

$$(3.8) \quad \ker D_A = \{0\}, \quad \text{ran } D_A \cap (\mathfrak{H} \ominus \mathfrak{H}_N^c) = \{0\}, \\ \text{and} \quad \text{ran } D_A \cap (\mathfrak{H} \ominus \mathfrak{H}_N^o) = \{0\}.$$

*In particular, if  $\ker D_A = \{0\}$  and  $\mathfrak{H} = \mathfrak{H}_N^c = \mathfrak{H}_N^o$ , then  $\tau$  is minimal; if  $\tau$  is minimal,  $\mathfrak{H}_N^c$  is invariant under  $A^*$ , and  $\mathfrak{H}_N^o$  is invariant under  $A$ , then  $\ker D_A = \{0\}$  and  $\mathfrak{H} = \mathfrak{H}_N^c = \mathfrak{H}_N^o$ .*

*Proof.* (i) Assume that (3.5) holds. Let  $f \in (\mathfrak{H}^c)^\perp$ . It follows from (3.2) and (3.4) that  $D_A f \in \bigcap_{n=0}^{\infty} \ker (M^* A^{*n}) = (\mathfrak{H}_N^c)^\perp$ . The second condition in (3.5) shows that  $D_A f = 0$  and the first condition in (3.5) yields  $f = 0$ . Therefore,  $\mathfrak{H}^c = \mathfrak{H}$  and  $\tau$  is controllable.

Now assume that  $\tau$  is controllable, i.e.  $\mathfrak{H}^c = \mathfrak{H}$ . Then (3.2) implies that  $\ker D_A = \{0\}$ . Furthermore, if  $D_A f \in (\mathfrak{H}_N^c)^\perp$ , then (3.4) implies that  $D_A f \in \bigcap_{n=0}^{\infty} \ker (M^* A^{*n})$ .

By (3.2) this leads to  $f \in (\mathfrak{H}^c)^\perp$  and hence  $f = 0$  by controllability of  $\tau$ . This shows that (3.5) is satisfied.

If  $\ker D_A = \{0\}$  and  $\mathfrak{H}_N^c = \mathfrak{H}$ , then (3.5) is satisfied. It follows that  $\tau$  is controllable.

If  $\mathfrak{H}_N^c$  is invariant under  $A^*$ , then  $\mathfrak{H} \ominus \mathfrak{H}_N^c \subset \mathfrak{H} \ominus \mathfrak{H}^c$  by Lemma 3.1. Hence, if in addition,  $\tau$  is controllable, it follows that  $\mathfrak{H}_N^c = \mathfrak{H}$ ; moreover, it follows that  $\ker D_A = \{0\}$ .

(ii) The proof is completely analogous to the proof for part (i).

(iii) If  $\tau$  is simple then it immediately follows from (3.1) that  $\ker D_A = \{0\}$ . Moreover, it is clear from (3.2) and (3.4) that  $D_A f \in \mathfrak{H} \ominus (\mathfrak{H}_N^c + \mathfrak{H}_N^o)$  if and only if  $f \in \mathfrak{H} \ominus (\mathfrak{H}^c + \mathfrak{H}^o)$ . Now the statement is obtained as in part (i).

(iv) This is obvious from the definition of minimality.  $\square$

**Corollary 3.3.** *Let the main operator  $A$  of the passive system  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be normal and let the system be simple. Then the system  $\tau$  is strongly stable and strongly co-stable.*

*Proof.* Since  $\tau$  is simple and  $A$  is normal, Proposition 3.2 shows that  $\ker D_A = \{0\}$  or, equivalently, that the contraction  $A$  is strict. Hence Lemma 2.2 implies that  $A \in C_{00}$ . Therefore  $\tau$  is strongly stable and strongly co-stable.  $\square$

**3.2. Defect functions.** Associated with  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  are the right and left *defect functions* (or *spectral factors*)  $\varphi_\Theta(\lambda)$  and  $\psi_\Theta(\lambda)$ , which satisfy

$$(3.9) \quad \varphi_\Theta^*(\xi)\varphi_\Theta(\xi) \leq I_{\mathfrak{M}} - \Theta^*(\xi)\Theta(\xi), \quad \psi_\Theta(\xi)\psi_\Theta^*(\xi) \leq I_{\mathfrak{N}} - \Theta(\xi)\Theta^*(\xi),$$

almost everywhere on  $\mathbb{T}$ . These operator-valued Schur functions are (up to a constant unitary factor) uniquely determined by the following maximality property: if  $\tilde{\varphi}(\lambda)$  and  $\tilde{\psi}(\lambda)$  are operator-valued Schur functions for which

$$(3.10) \quad \tilde{\varphi}^*(\xi)\tilde{\varphi}(\xi) \leq I_{\mathfrak{M}} - \Theta^*(\xi)\Theta(\xi), \quad \tilde{\psi}(\xi)\tilde{\psi}^*(\xi) \leq I_{\mathfrak{N}} - \Theta(\xi)\Theta^*(\xi),$$

then they are dominated by  $\varphi_\Theta(\lambda)$  and  $\psi_\Theta(\lambda)$  in the following sense:

$$(3.11) \quad \tilde{\varphi}^*(\xi)\tilde{\varphi}(\xi) \leq \varphi_\Theta^*(\xi)\varphi_\Theta(\xi), \quad \tilde{\psi}(\xi)\tilde{\psi}^*(\xi) \leq \psi_\Theta(\xi)\psi_\Theta^*(\xi),$$

almost everywhere on the unit circle  $\mathbb{T}$ ; cf. [17], [19], [20], [21], [22].

Note that it follows from Theorem 2.7 that the functions  $\varphi(\lambda)$  and  $\psi(\lambda)$  satisfy the inequalities

$$(3.12) \quad \varphi(\xi)^*\varphi(\xi) \leq \varphi_\Theta^*(\xi)\varphi_\Theta(\xi), \quad \psi(\xi)\psi^*(\xi) \leq \psi_\Theta(\xi)\psi_\Theta^*(\xi),$$

for almost all  $\xi \in \mathbb{T}$ .

**Proposition 3.4.** *Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system with a normal main operator  $A$  and let  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  be its transfer function. If  $\tau$  is simple and  $\varphi_\Theta(\lambda) = 0$  ( $\psi_\Theta(\lambda) = 0$ ), then  $\Theta(\lambda)$  is inner (co-inner, respectively).*

*Proof.* By Corollary 3.3 one has  $A \in C_{00}$  and, in particular,  $A$  is completely non-unitary. Therefore,  $\Phi_A(\lambda)$  and  $\Phi_{A^*}(\lambda)$  are bi-inner. On the other hand, if  $\varphi_\Theta(\lambda) = 0$  ( $\psi_\Theta(\lambda) = 0$ ), then (3.12) shows that  $\varphi(\xi) = 0$  ( $\psi(\xi) = 0$ ) for almost all  $\xi \in \mathbb{T}$ . Now (2.23) ((2.24), respectively) yields that  $D_{\Theta(\xi)} = 0$  ( $D_{\Theta^*(\xi)} = 0$ ) almost everywhere on  $\mathbb{T}$ , i.e.  $\Theta(\lambda)$  is inner (co-inner).  $\square$

**3.3. Unitary similarity.** Recall that two passive systems  $\tau_j = \{T_j; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_j\}$ ,  $j = 1, 2$ , are said to be unitarily similar if there is a unitary operator  $U : \mathfrak{H}_1 \rightarrow \mathfrak{H}_2$ , such that (1.4) holds. In particular, in this case the spectra of the corresponding main operators  $A_1$  and  $A_2$  coincide. It is clear that if the systems  $\tau_1$  and  $\tau_2$  are unitarily similar then they have the same transfer function. However, two minimal passive systems  $\tau_1$  and  $\tau_2$  with the same transfer function  $\Theta(\lambda)$  are in general not unitarily similar; such systems are only weakly similar as shown in D.Z. Arov [10], see (1.6). In the case of passive systems with normal main operators the following sufficient spectral-theoretic condition can be established.

**Theorem 3.5.** *Let  $\tau_1 = \{T_1; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_1\}$  and  $\tau_2 = \{T_2; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_2\}$  be two minimal passive systems whose transfer functions coincide in some neighborhood of zero. Let the main operator  $A_k$  be normal and let  $C_k = SB_k^*$ ,  $k = 1, 2$ , with  $S$  bounded and injective. Then, if the spectrum  $\sigma(A_k)$  of  $A_k$ ,  $k = 1, 2$ , does not contain interior points and  $\rho(A_1) \cap \rho(A_2)$  is a connected set in  $\mathbb{C}$ , the systems  $\tau_1$  and  $\tau_2$  are unitarily similar.*

*Proof.* Assume that the transfer functions  $\Theta_1(\lambda)$  and  $\Theta_2(\lambda)$  of  $\tau_1$  and  $\tau_2$  coincide in some neighborhood of zero. Since  $\Theta_1(\lambda)$  and  $\Theta_2(\lambda)$  are holomorphic on  $\mathbb{D}$  it follows that  $\Theta_1(\lambda) = \Theta_2(\lambda)$  for all  $\lambda \in \mathbb{D}$ . The definition (1.5) implies that  $D_1 = \Theta_1(0) = \Theta_2(0) = D_2$  and that

$$\sum_{m=0}^{\infty} \lambda^m C_1 A_1^m B_1 = \sum_{m=0}^{\infty} \lambda^m C_2 A_2^m B_2, \quad \lambda \in \mathbb{D}.$$

Since  $C_k = SB_k^*$ ,  $k = 1, 2$ , where  $S$  is bounded and injective, the previous equality yields

$$(3.13) \quad B_1^* A_1^m B_1 = B_2^* A_2^m B_2, \quad m \in \mathbb{N}_0.$$

Now define the relation  $Z_0$  by

$$(3.14) \quad Z_0 = \left\{ \left\{ \sum_{j=0}^m A_1^j B_1 u_j, \sum_{j=0}^m A_2^j B_2 u_j \right\} : u_0, u_1, \dots, u_m \in \mathfrak{M}, m \in \mathbb{N}_0 \right\}.$$

Clearly  $Z_0$  is linear and

$$\text{dom } Z_0 = \text{span} \{ A_1^n B_1 \mathfrak{M} : n \in \mathbb{N}_0 \}, \quad \text{ran } Z_0 = \text{span} \{ A_2^n B_2 \mathfrak{M} : n \in \mathbb{N}_0 \}.$$

Furthermore, it follows from (3.13) that

$$(3.15) \quad \left\{ \sum_{j=0}^m A_2^{*j} B_2 v_j, \sum_{j=0}^m A_1^{*j} B_1 v_j \right\} \in Z_0^*, \quad v_1, \dots, v_m \in \mathfrak{M}, \quad m \in \mathbb{N}_0,$$

so that

$$\text{span} \{ A_2^{*n} B_2 \mathfrak{M} : n \in \mathbb{N}_0 \} \subset \text{dom } Z_0^*, \quad \text{span} \{ A_1^{*n} B_1 \mathfrak{M} : n \in \mathbb{N}_0 \} \subset \text{ran } Z_0^*.$$

Due to the controllability and observability conditions (note that  $C_k^* = B_k S^*$ ), it follows from (3.14) and (3.15) that both  $Z_0$  and  $Z_0^*$  have dense domains and dense ranges. In particular,  $Z_0$  and  $Z_0^*$  are (graphs of) operators, and, in fact,  $\text{mul } Z_0^{**} = (\text{dom } Z_0^*)^\perp$  implies that  $Z_0$  is a closable operator, i.e., its closure  $Z_0^{**}$  is (the graph of) an operator; cf. [11].

Next it is shown that under the assumptions on the main operators  $A_k$ ,  $k = 1, 2$ , the mapping  $Z_0$  becomes isometric. Since  $A$  is contractive the spectrum  $\sigma(A_k)$  is a compact subset of the closed unit disk. The union  $\sigma(A_1) \cup \sigma(A_2)$  is also compact and, in addition, does not have interior points. Indeed, this follows immediately from the fact that the sets  $\sigma(A_1)$  and  $\sigma(A_2)$  are closed and do not have interior points. Furthermore, by assumption  $\mathbb{C} \setminus (\sigma(A_1) \cup \sigma(A_2)) = \rho(A_1) \cap \rho(A_2)$  is connected. Therefore, according to Mergelyan's theorem (see e.g. [26, Theorem 20.5]) every continuous complex-valued function on  $\sigma(A_1) \cup \sigma(A_2)$  can be uniformly approximated on  $\sigma(A_1) \cup \sigma(A_2)$  by complex polynomials. Since for every  $n, m \in \mathbb{N}_0$  the function  $f_{n,m}(z) = \bar{z}^n z^m$  is continuous on  $\mathbb{C}$ , there exists a sequence  $\{P_j^{n,m}(z) : j \in \mathbb{N}_0\}$  of polynomials converging uniformly on  $\sigma(A_1) \cup \sigma(A_2)$  to  $f_{n,m}(z)$ . It follows from (3.13) that for every  $n, k, j \in \mathbb{N}_0$  one has

$$(3.16) \quad B_1^* P_j^{n,m}(A_1) B_1 = B_2^* P_j^{n,m}(A_2) B_2.$$

The functional calculus for normal operators shows that  $f_{n,m}(A_k) = A_k^{*n} A_k^m$ ,  $k = 1, 2$ , and therefore taking strong limits in (3.16) yields

$$(3.17) \quad B_1^* A_1^{*n} A_1^m B_1 = B_2^* A_2^{*n} A_2^m B_2, \quad m, n \in \mathbb{N}_0.$$

These identities imply that

$$\left\| \sum_{j=0}^m A_1^j B_1 u_j \right\|^2 = \left\| \sum_{j=0}^m A_2^j B_2 u_j \right\|^2, \quad u_0, u_1, \dots, u_m \in \mathfrak{M}, \quad m \in \mathbb{N}_0,$$

and, therefore, the operator  $Z_0$  in (3.14) is isometric. Since  $Z_0$  is densely defined with dense range, its closure  $Z$  is unitary. The identities  $ZA_1 = A_2Z$  and  $ZB_1 = B_2$  are immediate from (3.14), while (3.15) shows that  $Z_0^* B_2 = B_1$  which gives the identity  $C_2 Z = C_1$ . Therefore, the systems  $\tau_1$  and  $\tau_2$  are unitarily similar; cf. (1.4).  $\square$

**Corollary 3.6.** *Let  $\tau_1 = \{T_1; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_1\}$  and  $\tau_2 = \{T_2; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_2\}$  be two minimal passive systems such that  $A_k$  is selfadjoint ( $A_k = A_k^*$ ) or skew-symmetric ( $A_k = -A_k^*$ ) and  $C_k = SB_k^*$ ,  $k = 1, 2$ , with  $S$  bounded and injective. Then  $\tau_1$  and  $\tau_2$  are unitarily similar if and only if their transfer functions coincide in some neighborhood of zero.*

**Corollary 3.7.** *Let  $\tau_1 = \{T_1; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_1\}$  and  $\tau_2 = \{T_2; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_2\}$  be two minimal passive systems such that  $A_k$  is normal and has a discrete spectrum, and  $C_k = SB_k^*$ ,  $k = 1, 2$ , with  $S$  bounded and injective. Then  $\tau_1$  and  $\tau_2$  are unitarily similar if and only if their transfer functions coincide in some neighborhood of zero.*

**Corollary 3.8.** *Let  $\tau_1 = \{T_1; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_1\}$  and  $\tau_2 = \{T_2; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_2\}$  be two minimal passive systems with a finite-dimensional state space  $\mathfrak{H}_k$  such that  $A_k$  is normal and  $C_k = SB_k^*$ ,  $k = 1, 2$ , with  $S$  bounded and injective. Then  $\tau_1$  and  $\tau_2$  are unitarily similar if and only if their transfer functions coincide in some neighborhood of zero.*

**Remark 3.9.** (i) The proof of Theorem 3.5 uses that fact that the operators  $f_{n,m}(A) = A^{*n} A^m$ ,  $m, n \in \mathbb{N}_0$ , can be approximated by a sequence of polynomials in  $A$ . If, in particular, the adjoint  $A^*$  of a bounded operator  $A$  can be approximated by a sequence  $P_n(A)$ ,  $n \in \mathbb{N}_0$ , of polynomials in  $A$ , i.e.,

$$(3.18) \quad A^* = s - \lim_{n \rightarrow \infty} P_n(A),$$

then the same is true for all of the operators  $f_{n,m}(A) = A^{*n}A^m$ ,  $m, n \in \mathbb{N}_0$ . By taking strong limits in  $AP_n(A) = P_n(A)A$  one obtains from (3.18) the identity  $AA^* = A^*A$ . Therefore, the condition (3.18) implies that  $A$  is a normal operator.

(ii) If  $A$  is a normal operator, then  $A^* = f(A)$  with  $f(z) = \bar{z}$  by the functional calculus for normal operators. The function  $f(z)$  does not satisfy the Cauchy-Riemann equations, so it is nowhere holomorphic. Consequently, if  $\sigma(A)$  has interior points, the adjoint  $A^*$  cannot satisfy the condition (3.18), as one would get a uniform approximation for  $f(z)$  on  $\sigma(A)$  via polynomials  $P_n(z)$ .

(iii) If  $A$  is a normal operator on a finite-dimensional space, then it has  $n = \dim \mathfrak{H}$  eigenvalues and it is unitarily similar to a diagonal matrix. Therefore, if  $A$  has  $d$  nonreal eigenvalues then by standard interpolation one finds a polynomial  $Q$  (say, of degree at most  $d - 1$  when using only the nonreal spectral points) such that  $A^* = Q(A)$ . If there are two normal operators  $A_1$  and  $A_2$  on  $\mathfrak{H}_k$ ,  $n_k = \dim \mathfrak{H}_k < \infty$ ,  $k = 1, 2$ , then together they have at most  $n_1 + n_2$  different nonreal eigenvalues and one can find a polynomial  $P$  (of degree at most  $n_1 + n_2 - 1$ ) such that  $A_1^* = P(A_1)$  and  $A_2^* = P(A_2)$ . Then  $f_{n,m}(A_k) = A_k^{*n}A_k^m = P(A_k)^n A_k^m$ ,  $n, m \in \mathbb{N}_0$ , is also a polynomial in  $A_k$ ,  $k = 1, 2$ . So, in the proof of Theorem 3.5 no limit procedure is needed in the case of finite-dimensional state spaces.

(iv) Finally, note that the criterion for unitary similarity of minimal passive systems with the same transfer function which has been established in [14] is essentially of different nature than the above spectral theoretical sufficient condition in Theorem 3.5.

#### 4. PASSIVE QUASI-SELFADJOINT SYSTEMS

**4.1. Quasi-selfadjoint contractions and associated passive systems.** Let  $\mathcal{H}$  be a Hilbert space. A linear operator  $T \in \mathbf{L}(\mathcal{H})$  is said to be a *quasi-selfadjoint contraction* (*qsc-operator* for short) if

$$\text{dom } T = \mathcal{H}, \quad \|T\| \leq 1, \quad \text{and } \ker (T - T^*) \neq \{0\}.$$

The next theorem is a consequence of Theorem 2.5; see [4].

**Theorem 4.1.** *Let  $T$  be a qsc-operator in the Hilbert space  $\mathcal{H}$  and let  $\mathfrak{N}$  be a subspace in  $\mathcal{H}$  such that  $\text{ran } (T - T^*) \subset \mathfrak{N}$ . Then with respect to the decomposition  $\mathcal{H} = \mathfrak{N} \oplus \mathfrak{H}$ , where  $\mathfrak{H} = \mathcal{H} \ominus \mathfrak{N}$ , the operator  $T$  has the following block form*

$$(4.1) \quad T = \begin{pmatrix} -KAK^* + D_{K^*}XD_{K^*} & KD_A \\ D_AK^* & A \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix},$$

where  $A = P_{\mathfrak{H}}T|_{\mathfrak{H}}$  is a selfadjoint contraction and  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$ ,  $X \in \mathbf{L}(\mathfrak{D}_{K^*})$  are contractions.

The system  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  is said to be *passive quasi-selfadjoint* ( $\tau$  is a *pqs-system* for short) if  $T$  in (1.7) is a contraction and if  $\text{ran } (T - T^*) \subset \mathfrak{N}$ . It follows that  $T$  is a qsc-operator in  $\mathfrak{N} \oplus \mathfrak{H}$  and that  $A = A^*$  and  $C = B^*$ . Moreover, according to Theorem 4.1,  $B$ ,  $C$ , and  $D$  have the form

$$(4.2) \quad B = D_AK^*, \quad C = KD_A, \quad D = -KAK^* + D_{K^*}XD_{K^*},$$

where  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$  and  $X \in \mathbf{L}(\mathfrak{D}_K^*)$  are contractions. For a  $pqs$ -system the controllable and observable subspaces coincide, see (3.1):

$$(4.3) \quad \mathfrak{H}^c = \mathfrak{H}^o = \overline{\text{span}} \{ D_A A^n K^* \mathfrak{N} : n \in \mathbb{N}_0 \} \subset \mathfrak{D}_A.$$

**4.2. Minimal representations of  $pqs$ -systems and unitary similarity.** A  $pqs$ -system can always be reduced to a minimal  $pqs$ -system.

**Proposition 4.2.** *Let  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  be a  $pqs$ -system of the form (1.7) and let  $B, C$ , and  $D$  be given by (4.2) with some contractions  $K$  and  $X$ . Define the system*

$$(4.4) \quad \tau_s = \{T_s; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}^s\},$$

where the subspace  $\mathfrak{H}^s$  is given by

$$(4.5) \quad \mathfrak{H}^s = \overline{\text{span}} \{ A^n K^* \mathfrak{N} : n \in \mathbb{N}_0 \},$$

and where the operator  $T_s$  is given by

$$(4.6) \quad T_s = \begin{pmatrix} D & C|_{\mathfrak{H}^s} \\ B & A|_{\mathfrak{H}^s} \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H}^s \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H}^s \end{pmatrix}.$$

Then  $\tau_s$  is a minimal  $pqs$ -system and the transfer functions of the systems  $\tau$  and  $\tau^s$  coincide. Moreover, the system  $\tau$  is minimal if and only if

- (i)  $\|Af\| < \|f\|$  for all  $f \in \mathfrak{H} \setminus \{0\}$ ,
- (ii)  $\mathfrak{H}^s = \mathfrak{H}$ .

In this case the system  $\tau$  is strongly stable and strongly co-stable.

*Proof.* The subspace  $\mathfrak{H}^s$  in (4.5) reduces  $A$  and therefore it also reduces  $D_A = (I_{\mathfrak{H}} - A^2)^{1/2}$ . Furthermore,  $\overline{\text{ran}} K^* \subset \mathfrak{H}^s$ . Let  $A_s = A|_{\mathfrak{H}^s}$ , then  $D_{A_s} = D_A|_{\mathfrak{H}^s}$  and, hence,  $D_A K^* = D_{A_s} K^*$ . Define the operator  $C_s$  by

$$C_s = C|_{\mathfrak{H}^s} = K D_{A_s}.$$

Then  $T_s$  in (4.6) is a  $qsc$ -operator in  $\mathfrak{N} \oplus \mathfrak{H}^s$ . Since  $\text{ran } K^* \subset \mathfrak{D}_A \cap \mathfrak{H}^s$ , one has  $\mathfrak{D}_{A_s} = \mathfrak{H}^s$ . Now the construction shows that the system  $\tau^s$  in (4.4) is minimal. Clearly, the transfer functions of  $\tau$  and  $\tau^s$  coincide.

As to the minimality of  $\tau$  observe that  $\mathfrak{H}^s = \mathfrak{H}_N^c = \mathfrak{H}_N^o$ , since  $A = A^*$ ; see (3.3). Hence, the characteristic properties (i) and (ii) for minimality of a  $pqs$ -system  $\tau$  are obtained from Proposition 3.2.

The last statement holds by Corollary 3.3. □

It is a consequence of Theorem 3.5 that within the class of  $pqs$ -systems the following unitary similarity criterion holds; see Corollary 3.6.

**Proposition 4.3.** *Let  $\tau_1 = \{T_1; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_1\}$  and  $\tau_2 = \{T_2; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}_2\}$  be two minimal  $pqs$ -systems. Then  $\tau_1$  and  $\tau_2$  are unitarily similar if and only if their transfer functions coincide in some neighborhood of zero.*

**4.3. Transfer functions of  $pqs$ -systems.** Let  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  be a  $pqs$ -system of the form (1.7) and assume that  $T$  is represented in the form (4.1). Then the transfer function  $\Theta(\lambda)$  of  $\tau$  has the form

$$(4.7) \quad \Theta(\lambda) = K\Phi_A(\lambda)K^* + D_{K^*}XD_{K^*}, \quad \lambda \in \mathbb{D},$$

where  $\Phi_A(\lambda)$  is the characteristic function of the selfadjoint contraction  $A$ ; see (2.6). The function  $\Phi_A(\lambda)$  is holomorphic on  $\mathbb{T} \setminus \{-1, 1\}$  and, in fact, it belongs to Herglotz-Neuman class on  $\text{Ext}\{(-\infty, -1] \cup [1, \infty)\}$ . Furthermore,  $\Phi_A(\lambda)$  has nontangential strong limit values  $\Phi_A(\pm 1) = \pm I_{\mathfrak{D}_A}$ ; see e.g. [4, Theorem 2.3]. Consequently, the limit value  $\Phi_A(\xi)$  is unitary for every  $\xi \in \mathbb{T}$  (see (2.7), (2.11)), in particular,  $\Phi_A(\lambda)$  is bi-inner. It follows from (4.7) that  $\Theta(\lambda)$ , initially defined on  $\mathbb{D}$ , admits a holomorphic continuation onto  $\text{Ext}\{(-\infty, -1] \cup [1, \infty)\}$ . Furthermore,  $\Theta(\lambda)$  has nontangential strong limit values  $\Theta(\pm 1)$  at  $\pm 1$  which are given by

$$(4.8) \quad \Theta(1) = KK^* + D_{K^*}XD_{K^*}, \quad \Theta(-1) = -KK^* + D_{K^*}XD_{K^*}.$$

Define the function  $W(\lambda)$  by

$$(4.9) \quad W(\lambda) = \Theta(\lambda) - \Theta(0), \quad \lambda \in \text{Ext}\{(-\infty, -1] \cup [1, \infty)\}.$$

Since

$$(4.10) \quad \Theta(0) = -KAK^* + D_{K^*}XD_{K^*},$$

it follows that

$$(4.11) \quad W(\lambda) = \lambda K(I - \lambda A)^{-1}D_A^2K^*, \quad \lambda \in \text{Ext}\{(-\infty, -1] \cup [1, \infty)\}.$$

Hence  $W^*(\bar{\lambda}) = W(\lambda)$  and

$$\frac{W(\lambda) - W^*(\xi)}{\lambda - \bar{\xi}} = \begin{cases} KD_A(I - \lambda A)^{-1}(I - \bar{\xi}A)^{-1}D_AK^*, & \bar{\xi} \neq \lambda, \\ KD_A(I - \lambda A)^{-2}D_AK^*, & \bar{\xi} = \lambda. \end{cases}$$

Therefore  $W(\lambda)$  is an operator-valued Herglotz-Neuman function with a holomorphic continuation onto  $\text{Ext}\{(-\infty, -1] \cup [1, \infty)\}$ . From (4.8) one sees that the strong limit values  $W(\pm 1)$  exist and that they are given by

$$W(1) = K(I + A)K^*, \quad W(-1) = -K(I - A)K^*.$$

Hence,

$$\frac{W(1) + W(-1)}{2} = KAK^*, \quad I - \frac{W(1) - W(-1)}{2} = I - KK^* = D_{K^*}^2 \geq 0.$$

Since  $X$  in (4.10) is a contraction in  $\mathfrak{D}_{K^*}$ , these identities show that

$$(4.12) \quad \Theta(0) \in \mathbf{B}\left(-\frac{W(1) + W(-1)}{2}, I - \frac{W(1) - W(-1)}{2}\right).$$

Here  $\mathbf{B}(S, R) = \{S + R^{1/2}XR^{1/2} \in \mathbf{L}(\mathfrak{N}) : X \text{ a contraction in } \mathbf{L}(\overline{\text{ran}} R)\}$  stands for the operator ball with center  $S \in \mathbf{L}(\mathfrak{N})$  and left and right radii  $R \geq 0$ .

**Definition 4.4.** Let  $\mathfrak{N}$  be a Hilbert space. The class  $\mathbf{S}^{qs}(\mathfrak{N})$  consists of all  $\mathbf{L}(\mathfrak{N})$ -valued functions  $\Theta(\lambda)$ , defined on  $\mathbb{D}$ , such that

- (S1)  $W(\lambda) = \Theta(\lambda) - \Theta(0)$  is a Herglotz-Neuman function with a holomorphic continuation onto the domain  $\text{Ext}\{(-\infty, -1] \cup [1, \infty)\}$ ;

- (S2) the strong limit values  $W(\pm 1)$  exist and  $W(1) - W(-1) \leq 2I$ ;  
 (S3)  $\Theta(0)$  belongs to the operator ball in (4.12).

The following proposition is now clear.

**Proposition 4.5.** *Let  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  be a  $pqs$ -system. Then its transfer function  $\Theta(\lambda)$  belongs to  $\mathbf{S}^{qs}(\mathfrak{N})$ .*

## 5. THE CLASS $\mathbf{S}^{qs}$ AND ITS REALIZATION VIA PASSIVE SYSTEMS

**5.1. The realization of the class  $\mathbf{S}^{qs}$ .** The next theorem is a converse to Proposition 4.5. In its proof a minimal  $pqs$ -system is constructed explicitly via an operator representation of the Herglotz-Nevanlinna function  $W(\lambda) = \Theta(\lambda) - \Theta(0)$ .

**Theorem 5.1.** *Let  $\mathfrak{N}$  be a Hilbert space and let  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ . Then  $\Theta(\lambda)$  is the transfer function of a minimal  $pqs$ -system  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$ .*

*Proof.* Assume that  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ . By the condition (S1) the function

$$\widetilde{W}(z) := -W(1/z), \quad z \in \text{Ext}[-1, 1],$$

is a Herglotz-Nevanlinna function of the class  $\mathbf{N}_{\mathfrak{N}}[-1, 1]$  with  $\widetilde{W}(\infty) = 0$ , see [4]. It follows from the condition (S2) that the strong limit values  $\widetilde{W}(\pm 1)$  exist. Then according to [4, Theorem 2.3] there exist a Hilbert space  $\widetilde{\mathfrak{H}}$ , a selfadjoint contraction  $\widetilde{A}$  in  $\widetilde{\mathfrak{H}}$ , and an operator  $\widetilde{G} \in \mathbf{L}(\mathfrak{N}, \mathfrak{D}_{\widetilde{A}})$ , such that

$$\widetilde{W}(z) = \widetilde{G}^*(\widetilde{A} - zI)^{-1}(I - \widetilde{A}^2)\widetilde{G},$$

see [4]. It follows that

$$W(-1) = -\widetilde{W}(-1) = -\widetilde{G}^*(I - \widetilde{A})\widetilde{G}, \quad W(1) = -\widetilde{W}(1) = \widetilde{G}^*(I + \widetilde{A})\widetilde{G}.$$

Consequently,

$$\frac{W(1) + W(-1)}{2} = \widetilde{G}^*\widetilde{A}\widetilde{G}, \quad I - \frac{W(1) - W(-1)}{2} = I - \widetilde{G}^*\widetilde{G}.$$

The condition  $W(1) - W(-1) \leq 2I$  implies that  $\widetilde{G}$  is contractive. The condition (S3) means that  $\Theta(0) = -\widetilde{G}^*\widetilde{A}\widetilde{G} + D_{\widetilde{G}}\widetilde{X}D_{\widetilde{G}}$  for some contraction  $\widetilde{X}$  in the Hilbert space  $\mathfrak{D}_{\widetilde{G}}$ . Define in the Hilbert space  $\mathcal{H} = \mathfrak{N} \oplus \widetilde{\mathfrak{H}}$  the operator  $\widetilde{T}$  by

$$\widetilde{T} = \begin{pmatrix} -\widetilde{G}^*\widetilde{A}\widetilde{G} + D_{\widetilde{G}}\widetilde{X}D_{\widetilde{G}} & \widetilde{G}^*D_{\widetilde{A}} \\ D_{\widetilde{A}}\widetilde{G} & \widetilde{A} \end{pmatrix}.$$

Then  $\widetilde{T}$  is a  $qsc$ -operator,  $\text{ran}(\widetilde{T} - \widetilde{T}^*) \subset \mathfrak{N}$ , and the operator  $\widetilde{T}$  defines a  $pqs$ -system  $\widetilde{\tau} = \{\widetilde{T}; \mathfrak{N}, \mathfrak{N}, \widetilde{\mathfrak{H}}\}$ ; cf. Theorem 4.1. The corresponding transfer function is given by

$$\Theta_{\widetilde{\tau}}(\lambda) = \widetilde{G}^* \left( -\widetilde{A} + \lambda \left( I - \lambda \widetilde{A} \right)^{-1} D_{\widetilde{A}}^2 \right) \widetilde{G} + D_{\widetilde{G}}\widetilde{X}D_{\widetilde{G}}, \quad \lambda \in \mathbb{D}.$$

Therefore,  $\Theta_{\widetilde{\tau}}(\lambda) = \Theta(0) + W(\lambda) = \Theta(\lambda)$ ,  $\lambda \in \mathbb{D}$ . This means that the function  $\Theta(\lambda)$  can be realized as the transfer function of the  $pqs$ -system  $\widetilde{\tau}$ . Finally, replacing  $\widetilde{\tau}$  by the system  $\widetilde{\tau}^s$ , cf. Proposition 4.2, one obtains a minimal  $pqs$ -system. The corresponding transfer function still coincides with the function  $\Theta(\lambda)$ .  $\square$

Observe that Theorem 5.1 implies that the class  $\mathbf{S}^{qs}(\mathfrak{N})$  is a subclass of the Schur class  $\mathbf{S}(\mathfrak{N})$ . Furthermore, the proof shows that a function  $\Theta(\lambda)$  from the class  $\mathbf{S}^{qs}(\mathfrak{N})$  admits the integral representation

$$\Theta(\lambda) = \Theta(0) + \lambda \int_{-1}^1 \frac{1-t^2}{1-t\lambda} d\Sigma(t),$$

where  $\Sigma(t)$  is a non-decreasing  $\mathbf{L}(\mathfrak{N})$ -valued function with bounded variation,  $\Sigma(-1) = 0$ ,  $\Sigma(1) \leq I_{\mathfrak{N}}$ , and

$$\left| \left( \left( \Theta(0) + \int_{-1}^1 t d\Sigma(t) \right) f, g \right) \right|^2 \leq ((I - \Sigma(1)) f, f) ((I - \Sigma(1)) g, g), \quad f, g \in \mathfrak{N}.$$

**Corollary 5.2.** *Let  $\mathfrak{N}$  be a Hilbert space and let  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ . If  $\varphi_{\Theta}(\lambda) = 0$  ( $\psi_{\Theta}(\lambda) = 0$ ) then  $\Theta(\lambda)$  is inner (co-inner).*

*Proof.* By Theorem 5.1 there exists a minimal  $pqs$ -system  $\tau = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  with transfer function  $\Theta(\lambda)$ . Now the statement follows from Proposition 3.4.  $\square$

**Theorem 5.3.** *Let  $\mathfrak{N}$  be a Hilbert space and let  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ . Then:*

(i) *if  $\Theta(\lambda)$  is inner then*

$$(5.1) \quad \left( \frac{\Theta(1) - \Theta(-1)}{2} \right)^2 = \frac{\Theta(1) - \Theta(-1)}{2},$$

$$(\Theta(1) + \Theta(-1))^* (\Theta(1) + \Theta(-1)) = 4I_{\mathfrak{N}} - 2(\Theta(1) - \Theta(-1));$$

(ii) *if  $\Theta(\lambda)$  is co-inner then*

$$(5.2) \quad \left( \frac{\Theta(1) - \Theta(-1)}{2} \right)^2 = \frac{\Theta(1) - \Theta(-1)}{2},$$

$$(\Theta(1) + \Theta(-1))(\Theta(1) + \Theta(-1))^* = 4I_{\mathfrak{N}} - 2(\Theta(1) - \Theta(-1));$$

(iii) *if (5.1) ((5.2)) holds and  $\Theta(\xi)$  is isometric (co-isometric) for some  $\xi \in \mathbb{T}$ ,  $\xi \neq \pm 1$ , then  $\Theta(\lambda)$  is inner (co-inner).*

*Proof.* Since  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ , it is the transfer function of a minimal  $pqs$ -system  $\tau = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$ . The operator  $T$ , being quasi-selfadjoint, has the form (4.1) and  $\Theta(\lambda)$  is given by (4.7) with a holomorphic continuation into the domain  $\text{Ext} \{(-\infty, -1] \cup [1, \infty)\}$ . Since  $\Phi_A(\xi)$  is unitary for every  $\xi \in \mathbb{T}$ , it follows from (2.23) and (2.24) in Theorem 2.7 and the definitions (2.20) and (2.21) that for all  $h \in \mathfrak{N}$  and  $\xi \in \mathbb{T}$

$$(5.3) \quad \|D_{\Theta(\xi)} h\|^2 = \|D_X D_{K^*} h\|^2 + \|(D_K \Phi_A(\xi) K^* - K^* X D_{K^*}) h\|^2,$$

$$\|D_{\Theta^*}(\xi) h\|^2 = \|D_{X^*} D_{K^*} h\|^2 + \|(D_K \Phi_A(\bar{\xi}) K^* - K^* X^* D_{K^*}) h\|^2.$$

(i) Suppose that  $\Theta(\lambda)$  is inner. Then (5.3) shows that

$$\begin{cases} D_X D_{K^*} = 0, \\ D_K \Phi_A(\xi) K^* = K^* X D_{K^*}, \quad \xi \in \mathbb{T}. \end{cases}$$

The last equality yields that  $D_K \Phi_A(\lambda) K^* = K^* X D_{K^*}$  for all  $\lambda \in \mathbb{D}$ . Since

$$\Phi_A(\lambda) = -A + \sum_{n=0}^{\infty} \lambda^{n+1} A^n D_A^2,$$

it follows that  $D_K D_A A^n D_A K^* = 0$ ,  $n \in \mathbb{N}_0$ . The minimality of  $\tau$  implies that  $D_K f = 0$  for all  $f \in \mathfrak{D}_A$ ; see (4.3). Hence,  $K$  is isometric and  $D_{K^*}$  is an orthogonal projector in the subspace  $\mathfrak{N}$ . Due to the identity  $D_X D_{K^*} = 0$ ,  $X \in \mathbf{L}(\mathfrak{D}_{K^*})$  is isometric (here possibly  $\mathfrak{D}_{K^*} = \{0\}$ ). Now from the equalities in (4.8) one obtains

$$(5.4) \quad KK^* = \frac{\Theta(1) - \Theta(-1)}{2}$$

and

$$(5.5) \quad XD_{K^*} = D_{K^*}XD_{K^*} = \frac{\Theta(1) + \Theta(-1)}{2}.$$

These identities together with the equality  $X^*X = I_{\mathfrak{D}_{K^*}}$  lead to (5.1).

(ii) The proof is similar to that of (i).

(iii) Assume that (5.1) holds and that  $\Theta(\xi)$  is isometric for some  $\xi \in \mathbb{T}$ ,  $\xi \neq \pm 1$ . Due to (5.1)  $D_{K^*}$  is an orthogonal projector in  $\mathfrak{N}$  and  $X$  is isometric in  $\mathfrak{D}_{K^*}$ . Hence,  $K^* \in \mathbf{L}(\mathfrak{N}, \mathfrak{H})$  is a partial isometry and, moreover,  $D_K K^* = K^* D_{K^*} = 0$  and  $K^* X D_{K^*} = 0$ . Since  $\Theta(\xi)$  is isometric, (5.3) gives  $D_K \Phi_A(\xi) K^* = 0$ . Furthermore, since

$$-A + \xi(I_{\mathfrak{H}} - A^2)(I_{\mathfrak{H}} - \xi A)^{-1} = \bar{\xi} + (\xi - \bar{\xi})(I_{\mathfrak{H}} - \xi A)^{-1},$$

the equality  $D_K \Phi_A(\xi) K^* = 0$  with  $\xi \neq \bar{\xi}$  implies that  $D_K(I_{\mathfrak{H}} - \xi A)^{-1} K^* = 0$ , i.e.

$$(I_{\mathfrak{H}} - \xi A)^{-1} \text{ran } K^* \subset \text{ran } K^*.$$

Thus  $A(\text{ran } K^*) \subset \text{ran } K^*$  (since  $A = A^*$ ) and hence  $D_A(\text{ran } K^*) \subset \text{ran } K^*$ , so that

$$(5.6) \quad \overline{\text{span}} \{A^n D_A K^* \mathfrak{N} : n \in \mathbb{N}_0\} \subset \text{ran } K^*.$$

Consequently  $\text{ran } K^* = \mathfrak{H}$ , i.e.,  $K \in \mathbf{L}(\mathfrak{H}, \mathfrak{N})$  is isometric. Therefore,  $D_K \Phi_A(\zeta) K^* = 0$  for all  $\zeta \in \mathbb{T}$  and  $\Theta(\lambda)$  is inner in view of (5.3). Similarly, if (5.2) holds and  $\Theta(\xi)$  is co-isometric for some  $\xi \in \mathbb{T}$ ,  $\xi \neq \pm 1$ , then  $\Theta(\lambda)$  is co-inner.  $\square$

**Theorem 5.4.** *Let the Hilbert space  $\mathfrak{N}$  be finite-dimensional and let  $\Theta(\lambda)$  be a non-constant inner function from  $\mathbf{S}^{qs}(\mathfrak{N})$ . Then  $\Theta(\lambda)$  is rational and*

$$\Theta(\lambda) = \text{diag} \left( \frac{\lambda - a_1}{1 - \lambda a_1}, \frac{\lambda - a_2}{1 - \lambda a_2}, \dots, \frac{\lambda - a_m}{1 - \lambda a_m}, X \right)$$

*relative to some orthonormal basis in  $\mathfrak{N}$ . Here the not necessarily distinct numbers  $a_1, a_2, \dots, a_m$  belong to  $(-1, 1)$ , and  $X$  is a constant unitary matrix.*

*Proof.* Since  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ , it is the transfer function of a minimal  $pqs$ -system  $\tau = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$ . As  $T$  is quasi-selfadjoint, it has the form (4.1), and  $\Theta(\lambda)$  is given by (4.7). Since  $\mathfrak{N}$  is finite-dimensional and  $\Theta(\lambda)$  is inner,  $\Theta(\lambda)$  is automatically bi-inner. Then in (3.9) one has  $\varphi_{\Theta}(\lambda) = 0$  and  $\psi_{\Theta}(\lambda) = 0$ . Thus by [5, Theorem 1.1]  $\tau$  is conservative (in fact, this conclusion can be derived also from the proof of Theorem 5.3 above by applying Corollary 2.6). As  $\Theta(\lambda)$  is nonconstant,  $A$  is non-isometric,  $K$  is isometric, and  $X$  appearing in (4.1) is unitary in  $\mathfrak{D}_{K^*}$ . Since  $\tau$  is minimal, Proposition 4.2 shows that  $\mathfrak{D}_A = \mathfrak{H} = \mathfrak{H}^s$ . Hence  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$  isometric implies that  $\dim \mathfrak{H} \leq \dim \mathfrak{N} < \infty$ , so that  $\Theta(\lambda)$  is rational. Let  $\mathfrak{N}_0 = \text{ran } K$ ,  $\mathfrak{N}_1 = \mathfrak{N} \ominus \mathfrak{N}_0 = \ker K^*$ . Then  $K^* \mathfrak{N}_0 = \mathfrak{H}$  and

$$\Theta(\lambda) \upharpoonright \mathfrak{N}_1 = X : \mathfrak{N}_1 \rightarrow \mathfrak{N}_1, \quad \Theta(\lambda) \upharpoonright \mathfrak{N}_0 = K \Phi_A(\lambda) K^* \upharpoonright \mathfrak{N}_0 : \mathfrak{N}_0 \rightarrow \mathfrak{N}_0.$$

Suppose that  $\dim \mathfrak{H} = m$  and that  $a_1, \dots, a_m$  are the eigenvalues of  $A$ . Choose an orthonormal basis in  $\mathfrak{H}$  consisting of eigenvectors of  $A$ . Then  $K$  maps this basis onto some orthonormal basis in  $\mathfrak{N}_0$ . With respect to this basis the matrix  $\Theta(\lambda)|_{\mathfrak{N}_0}$  is diagonal with entries  $\theta_{kk}(\lambda)$ ,  $k = 1, \dots, m$ , and since  $\Phi_A(\lambda)$  has the form (2.6), it follows that

$$\Theta_{kk}(\lambda) = -a_k + \frac{(1 - a_k^2)\lambda}{1 - \lambda a_k} = \frac{\lambda - a_k}{1 - \lambda a_k}.$$

This completes the proof.  $\square$

**5.2. Bi-inner dilations of functions from the class  $\mathbf{S}^{qs}$ .** The function  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  is said to have an *inner dilation* if there exists a function  $\Theta_r(\lambda)$  such that

$$\Theta(\lambda) = \begin{pmatrix} \Theta(\lambda) \\ \Theta_r(\lambda) \end{pmatrix} \in \mathbf{S}(\mathfrak{M}, \mathfrak{N} \oplus \mathfrak{L})$$

is inner. The function  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  is said to have a *co-inner dilation* if there exists a function  $\Theta_l(\lambda)$  such that

$$\Theta(\lambda) = \begin{pmatrix} \Theta(\lambda) & \Theta_l(\lambda) \end{pmatrix} \in \mathbf{S}(\mathfrak{M} \oplus \mathfrak{K}, \mathfrak{N})$$

is co-inner. The function  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$  is said to have a *bi-inner dilation* if there exist functions  $\Theta_{11}(\lambda)$ ,  $\Theta_{22}(\lambda)$ , and  $\Theta_{21}(\lambda)$  such that

$$(5.7) \quad \Theta(\lambda) = \begin{pmatrix} \Theta(\lambda) & \Theta_{12}(\lambda) \\ \Theta_{21}(\lambda) & \Theta_{22}(\lambda) \end{pmatrix} \in \mathbf{S}(\mathfrak{M} \oplus \mathfrak{K}, \mathfrak{N} \oplus \mathfrak{L})$$

is bi-inner. Recall the following result due to Arov [11]; cf. [16].

**Proposition 5.5.** *Let  $\tau = \{T; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}\}$  be a passive system with transfer function  $\Theta(\lambda)$ . Then:*

- (i) *if  $\tau$  is strongly stable, then  $\Theta(\lambda)$  has an inner dilation;*
- (ii) *if  $\tau$  is strongly co-stable, then  $\Theta(\lambda)$  has a co-inner dilation;*
- (iii) *if  $\tau$  is strongly stable and strongly co-stable, then  $\Theta(\lambda)$  has a bi-inner dilation.*

In [5] this result was proved using the parametrization in Theorem 2.5 and (2.22). Since a function  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  can be realized as the transfer function of a minimal  $pqs$ -system, it is strongly stable and strongly co-stable by Corollary 3.3. Hence it admits a bi-inner dilation by Proposition 5.5.

**Proposition 5.6.** *Among the bi-inner dilations of  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  there exists a bi-inner dilation from the class  $\mathbf{S}^{qs}(\mathfrak{N})$ .*

*Proof.* Since  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ , it is the transfer function of a minimal  $pqs$ -system  $\tau = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  by Theorem 5.1. The operator  $T$ , being quasi-selfadjoint, has the form (4.1) and therefore  $\Theta(\lambda)$  is given by (4.7). Define the following functions

$$\begin{aligned} \Theta_{21}(\lambda) &:= \begin{pmatrix} D_K \Phi_A(\lambda) K^* - K^* X D_{K^*} \\ -D_X D_{K^*} \end{pmatrix} : \mathfrak{N} \rightarrow \begin{pmatrix} \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix}, \quad \lambda \in \mathbb{D}, \\ \Theta_{12}(\lambda) &:= \begin{pmatrix} K \Phi_A(\lambda) D_K - D_{K^*} X K & D_{K^*} D_{X^*} \end{pmatrix} : \begin{pmatrix} \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix} \rightarrow \mathfrak{N}, \quad \lambda \in \mathbb{D}, \\ \Theta_{22}(\lambda) &= \begin{pmatrix} K^* X K + D_K \Phi_A(\lambda) D_K & -K^* D_{X^*} \\ D_X K & X^* \end{pmatrix} : \begin{pmatrix} \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix}, \quad \lambda \in \mathbb{D}. \end{aligned}$$

Let  $\mathfrak{K} = \mathfrak{L} = \mathfrak{D}_K \oplus \mathfrak{D}_{K^*}$  and  $\mathfrak{V} = \mathfrak{N} \oplus \mathfrak{L}$ , and let  $\Theta(\lambda)$  be defined by (5.7). Furthermore, define the operator  $\mathbf{T}$  by

$$\mathbf{T} = \begin{pmatrix} \mathbf{D} & \mathbf{C} \\ \mathbf{B} & \mathbf{A} \end{pmatrix} : \begin{pmatrix} \mathfrak{V} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{V} \\ \mathfrak{H} \end{pmatrix},$$

where  $\mathbf{A} = A$ ,  $\mathbf{D} = \Theta(0)$ , and

$$\mathbf{B} = D_A \begin{pmatrix} K^* & D_K & 0 \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix} \rightarrow \mathfrak{H}, \quad \mathbf{C} = \mathbf{B}^* = \begin{pmatrix} K \\ D_K \\ 0 \end{pmatrix} D_A : \mathfrak{H} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix}.$$

Simple calculations show that the operator  $\mathbf{T}$  is unitary and quasi-selfadjoint. Hence, the system  $\eta = \{\mathbf{T}; \mathfrak{V}, \mathfrak{V}, \mathfrak{H}\}$  is conservative. Since  $A \in C_{00}$ , the system  $\eta$  is minimal; see Corollary 2.4 and e.g. [5, Proposition 5.2]. In addition it is easy to see that the transfer function of  $\eta$  coincides with  $\Theta(\lambda)$  and therefore  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{V})$  by Proposition 4.5. Since  $\eta$  is minimal and conservative,  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{V})$  is a bi-inner dilation of  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{N})$  in view of [5, Corollary 5.3].  $\square$

**Remark 5.7.** With straightforward calculations it is easy to see directly that  $\Theta(\lambda) \in \mathbf{S}(\mathfrak{V})$  as defined explicitly in Proposition 5.6 is bi-inner; cf. [5, Proposition 7.1]. Furthermore, using the explicit formula of  $\Theta(\lambda)$  one can also calculate the function  $\mathbf{W}(\lambda) := \Theta(\lambda) - \Theta(0)$  as introduced in Definition 4.4. Observe, that

$$\mathbf{W}(\lambda) : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{D}_K \\ \mathfrak{D}_{K^*} \end{pmatrix}$$

and that

$$\Theta(0) = \begin{pmatrix} -KAK^* + D_{K^*}XD_{K^*} & -KAD_K - D_{K^*}XK & D_{K^*}D_{X^*} \\ -D_KAK^* - K^*XD_{K^*} & K^*XK - D_KAD_K & -K^*D_{X^*} \\ -D_XD_{K^*} & D_XK & X^* \end{pmatrix}.$$

Thus,

$$\mathbf{W}(\lambda) = \begin{pmatrix} \lambda KD_A^2(I_{\mathfrak{H}} - \lambda A)^{-1}K^* & \lambda KD_A^2(I_{\mathfrak{H}} - \lambda A)^{-1}D_K & 0 \\ \lambda D_KD_A^2(I_{\mathfrak{H}} - \lambda A)^{-1}K^* & \lambda D_KD_A^2(I_{\mathfrak{H}} - \lambda A)^{-1}D_K & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

which implies that the Nevanlinna kernel

$$\frac{\mathbf{W}(\lambda) - \mathbf{W}^*(\lambda)}{\lambda - \bar{\lambda}}$$

is given by

$$\begin{pmatrix} KD_A^2(I_{\mathfrak{H}} - \bar{\lambda}A)^{-1}(I_{\mathfrak{H}} - \lambda A)^{-1}K^* & KD_A^2(I_{\mathfrak{H}} - \bar{\lambda}A)^{-1}(I_{\mathfrak{H}} - \lambda A)^{-1}D_K & 0 \\ D_KD_A^2(I_{\mathfrak{H}} - \bar{\lambda}A)^{-1}(I_{\mathfrak{H}} - \lambda A)^{-1}K^* & D_KD_A^2(I_{\mathfrak{H}} - \bar{\lambda}A)^{-1}(I_{\mathfrak{H}} - \lambda A)^{-1}D_K & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore,  $\mathbf{W}(\lambda)$  is a Herglotz-Nevanlinna function defined on  $\text{Ext} \{(-\infty, -1] \cup [1, \infty)\}$ , which reflects the defining properties of the class  $\mathbf{S}^{qs}(\mathfrak{N})$  in Definition 4.4.

6. MINIMAL SYSTEMS WITH TRANSFER FUNCTIONS OF THE CLASS  $\mathbf{S}^{qs}$ 

In this section the class of minimal systems  $\dot{\tau} = \{T, \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  of the form (1.2) are considered. The system  $\dot{\tau}$  is not assumed to be passive, so that the bounded operator  $T$  in (1.1) need not be contractive. The next result can be seen as an extension of the unitary similarity result for  $pqs$ -systems in Proposition 4.3.

**Theorem 6.1.** *Let  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  and let*

$$(6.1) \quad \dot{\tau} = \left\{ \begin{pmatrix} \dot{D} & \dot{C} \\ \dot{B} & \dot{A} \end{pmatrix}; \mathfrak{N}, \mathfrak{N}, \mathfrak{H} \right\}.$$

*be a minimal, not necessarily passive, system whose transfer function coincides with  $\Theta(\lambda)$  in some neighborhood of zero. Then there exists a positive selfadjoint operator  $S$  in  $\mathfrak{H}$  such that*

$$\dot{A}^* S = S \dot{A}, \quad \dot{C}^* = S \dot{B},$$

*and such that*

$$\widehat{\tau}_0 = \left\{ \begin{pmatrix} D & \dot{C} S^{-1/2} \\ S^{1/2} \dot{B} & S^{1/2} \dot{A} S^{-1/2} \end{pmatrix}; \mathfrak{N}, \mathfrak{N}, \mathfrak{H} \right\}$$

*is a minimal  $pqs$ -system with transfer function  $\Theta(\lambda)$ . Moreover,  $\dot{\tau}$  becomes a  $pqs$ -system with respect to the inner product in  $\text{dom } S^{1/2} \subset \mathfrak{H}$  given by*

$$(\varphi, \psi)_{S^{1/2}} = (S^{1/2} \varphi, S^{1/2} \psi)_{\mathfrak{H}}, \quad \varphi, \psi \in \text{dom } S^{1/2}.$$

*Furthermore, if  $\Theta(\lambda)$  is inner (co-inner), then all minimal passive realizations of  $\Theta(\lambda)$  are unitarily similar isometric (co-isometric)  $pqs$ -systems.*

*Proof.* Since  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$ , it is the transfer function of a minimal  $pqs$ -system  $\tau_0$  of the form

$$(6.2) \quad \tau_0 = \left\{ \begin{pmatrix} D & C \\ B & A \end{pmatrix}; \mathfrak{N}, \mathfrak{N}, \overset{\circ}{\mathfrak{H}} \right\}.$$

Here  $A$  is a selfadjoint contraction in the Hilbert space  $\overset{\circ}{\mathfrak{H}}$  and  $C = B^*$ . Furthermore,  $\Theta(\lambda)$  has the form

$$\Theta(\lambda) = W(\lambda) + D,$$

where  $W(\lambda) = \lambda B^*(I - \lambda A)^{-1} B$ ,  $\lambda \in \mathbb{D}$ , is a Herglotz-Nevanlinna function on the domain  $\text{Ext } \{(-\infty, -1] \cup [1, \infty)\}$  and  $W(0) = 0$ . Since the transfer function of  $\dot{\tau}$  in (6.1) is of the form

$$\Theta(\lambda) = D + \lambda \dot{C}(I - \lambda \dot{A})^{-1} \dot{B},$$

one concludes that  $\dot{D} = D$  and in some neighborhood of zero

$$(6.3) \quad \dot{C}(I - \lambda \dot{A})^{-1} \dot{B} = B^*(I - \lambda A)^{-1} B.$$

Consequently,

$$(6.4) \quad \dot{C} \dot{A}^k \dot{B} = B^* A^k B, \quad k \in \mathbb{N}_0.$$

Define the linear operator  $Y$  by

$$(6.5) \quad Y = \left\{ \left\{ \sum_{k=0}^n \dot{A}^k \dot{B} u_k, \sum_{k=0}^n A^k B u_k \right\} : u_0, u_1, \dots, u_n \subset \mathfrak{N}, \quad n \in \mathbb{N}_0 \right\}.$$

It follows from (3.13) that

$$(6.6) \quad \left\{ \sum_{k=0}^m A^k B v_k, \sum_{k=0}^m \dot{A}^{*k} \dot{C}^* v_k \right\} \in Y^*, \quad v_1, \dots, v_m \in \mathfrak{N}, \quad m \in \mathbb{N}_0.$$

By minimality of  $\dot{\tau}$  and  $\tau_0$ ,  $Y$  and its adjoint  $Y^*$  have a dense domain and dense range. In particular, the operator  $Y$  is closable and the closure  $\bar{Y} = Y^{**}$  is a densely defined operator with dense range. Definition 6.5 and the relations (6.4), (6.6) yield the equalities

$$(6.7) \quad \begin{cases} A\bar{Y}u = \bar{Y}\dot{A}u, & u \in \text{dom } \bar{Y}, \\ \dot{C}u = B^*\bar{Y}u, & u \in \text{dom } \bar{Y}, \\ Y\dot{B} = B, \end{cases}$$

which means that the systems  $\dot{\tau}$  and  $\tau_0$  are weakly similar; see (1.6), cf. also [10]. Now define a positive selfadjoint operator  $S$  by  $S = Y^*\bar{Y}$ . Since  $Y^*B = \dot{C}^*$  and  $Y\dot{B} = B$ , one obtains  $S\dot{B} = \dot{C}^*$ . Represent  $\bar{Y}$  in the form  $\bar{Y} = US^{1/2}$ , where  $U : \mathfrak{H} \rightarrow \overset{0}{\mathfrak{H}}$  is an isometry and  $\text{dom } S^{1/2} = \text{dom } \bar{Y}$ . Because  $\overline{\text{ran}} Y = \overset{0}{\mathfrak{H}}$ , the operator  $U$  is a unitary from  $\mathfrak{H}$  onto  $\overset{0}{\mathfrak{H}}$ . Define a selfadjoint contraction  $\hat{A}$  in  $\mathfrak{H}$  by the equality:

$$\hat{A} = U^{-1}AU.$$

Then the equality  $\bar{Y}\dot{A} = A\bar{Y}$  gives

$$S^{1/2}\dot{A}v = \hat{A}S^{1/2}v, \quad v \in \text{dom } S^{1/2},$$

or, equivalently,

$$\hat{A}v = S^{1/2}\dot{A}S^{-1/2}v, \quad v \in \text{dom } S^{-1/2}.$$

Let  $\hat{B} = U^{-1}B$ . From  $Y\dot{B} = B$  one obtains  $S^{1/2}\dot{B} = \hat{B}$  and the relation  $S\dot{B} = \dot{C}^*$  yields  $S^{-1/2}\dot{C}^* = \hat{B}$ . Therefore  $\hat{B}^*v = \dot{C}S^{-1/2}v$ ,  $v \in \text{dom } S^{-1/2}$ . The system

$$\hat{\tau}_0 = \left\{ \begin{pmatrix} D & \hat{B}^* \\ \hat{B} & \hat{A} \end{pmatrix}; \mathfrak{N}, \mathfrak{N}, \mathfrak{H} \right\}$$

is unitarily equivalent to  $\tau_0$ . Hence  $\hat{\tau}_0$  is a minimal  $pqs$ -system with the transfer function  $\Theta(\lambda)$ . Let  $v \in \text{dom } S$ ,  $u \in \text{dom } S^{1/2}$ . Then

$$(6.8) \quad \begin{aligned} (S^{1/2}\dot{A}v, S^{1/2}u) &= (\hat{A}S^{1/2}v, S^{1/2}u) = (S^{1/2}v, \hat{A}S^{1/2}u) \\ &= (S^{1/2}v, S^{1/2}\dot{A}u) = (\dot{A}^*Sv, u). \end{aligned}$$

It follows that the vector  $S^{1/2}\dot{A}v$  belongs to  $\text{dom } S^{1/2}$  and

$$S\dot{A}v = \dot{A}^*Sv, \quad v \in \text{dom } S.$$

Consider  $\text{dom } S^{1/2}$  as a pre-Hilbert space equipped with the inner product  $(f, g)_{S^{1/2}} := (S^{1/2}f, S^{1/2}g)$  and let  $\mathfrak{H}_{S^{1/2}}$  be the completion of  $\text{dom } S^{1/2}$  with respect to this inner product. Since  $S^{1/2}\dot{A}v = \hat{A}S^{1/2}v$ ,  $v \in \text{dom } S^{1/2}$ , one obtains from (6.8)

$$(\dot{A}u, v)_{S^{1/2}} = (S^{1/2}\dot{A}v, S^{1/2}u) = (S^{1/2}v, S^{1/2}\dot{A}u) = (v, \dot{A}u)_{S^{1/2}},$$

for all  $v, u \in \text{dom } S^{1/2}$ . Thus, the operator  $\dot{A}$  is symmetric with respect to the inner product  $(\cdot, \cdot)_{S^{1/2}}$ . Furthermore, the operator  $\dot{T}$  defined via the block formula

$$\dot{T} = \begin{pmatrix} D & \dot{C} \\ \dot{B} & \dot{A} \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H}_{S^{1/2}} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H}_{S^{1/2}} \end{pmatrix}$$

is a *qsc*-operator. In fact, since  $\dot{C}^* = S^{1/2}\hat{B}$  and  $S^{1/2}\dot{B} = \hat{B}$ , the operator  $\dot{B} : \mathfrak{N} \rightarrow \mathfrak{H}_{S^{1/2}}$  is the adjoint of the operator  $\dot{C} : \text{dom } S^{1/2} \subset \mathfrak{H}_{S^{1/2}} \rightarrow \mathfrak{N}$ . Because the operator matrix

$$\hat{T} = \begin{pmatrix} D & \hat{B}^* \\ \hat{B} & \hat{A} \end{pmatrix} : \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{N} \\ \mathfrak{H} \end{pmatrix}$$

is a contraction, one obtains with  $f \in \text{dom } S^{1/2}$  and  $u \in \mathfrak{N}$

$$\begin{aligned} & \left\| \dot{A}f + \dot{B}u \right\|_{S^{1/2}}^2 + \left\| \dot{C}f + Du \right\|^2 \\ &= \left\| S^{1/2}(\dot{A}f + \dot{B}u) \right\|^2 + \left\| \dot{C}f + Du \right\|^2 \\ &= \left\| \hat{A}S^{1/2}f + \hat{B}u \right\|^2 + \left\| \hat{B}^*S^{1/2}f + Du \right\|^2 \\ &\leq \|S^{1/2}f\|^2 + \|u\|^2 = \|f\|_{S^{1/2}}^2 + \|u\|^2. \end{aligned}$$

Therefore, after a renormalization of the state space by means of the operator  $S$  the system  $\dot{\tau}$  becomes a *pqs*-system.

Finally, if the function  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  is inner (co-inner) then  $\varphi_\Theta(\xi) = 0$  ( $\psi_\Theta(\xi) = 0$ ) for almost all  $\xi \in \mathbb{T}$ , see (3.9), and now it follows from [5, Theorem 1.1] that every two minimal passive realizations of the inner (co-inner) function  $\Theta(\lambda) \in \mathbf{S}^{qs}(\mathfrak{N})$  are unitarily similar isometric (co-isometric) *pqs*-systems.  $\square$

## 7. THE CLASS $\mathbf{S}^{qs}$ AND $Q$ -FUNCTIONS OF QUASI-SELFADJOINT CONTRACTIONS

**7.1. Hermitian contractions.** Let  $A_0$  be a Hermitian contraction in the Hilbert space  $\mathcal{H}$  with  $\text{dom } A_0 = \mathfrak{H}$  and let  $\mathfrak{N} = \mathcal{H} \ominus \mathfrak{H}$ . Then  $A := P_{\mathfrak{H}}A_0|_{\mathfrak{H}}$  is selfadjoint in the Hilbert space  $\mathfrak{H}$ . It follows that there is a contraction  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$  such that

$$(7.1) \quad A_0 = A + KD_A.$$

The operator  $A_0$  is said to be *simple* if there is no nonzero subspace in  $\text{dom } A_0$  which is invariant under  $A_0$ ; cf. [24]. Since  $A_0$  is Hermitian, simplicity of  $A_0$  is equivalent to  $A_0$  being completely nonselfadjoint, i.e.,  $A_0$  has no selfadjoint part. Note that  $A_0$  is simple if and only if the subspace  $\mathfrak{H}_0^s$  in (4.5) coincides with  $\mathfrak{H}$ ; cf. [4, Lemma 3.2]. For a Hermitian contraction  $A_0 = A + KD_A$  the restriction  $A_0|_{\mathfrak{H}_0^s}$  is called the *simple part* of  $A_0$ .

A *qsc*-operator  $T$  is said to be a *quasi-selfadjoint contractive extension* or *qsc-extension* of a Hermitian contraction  $A_0$  if

$$A_0 \subset T \text{ and } A_0 \subset T^*,$$

or, equivalently, if  $\text{dom } A_0 \subset \ker(T - T^*)$ , cf. [6], [8]. A *qsc*-operator  $T$  has always Hermitian restrictions  $A_0$  for which  $T$  is a *qsc*-extension. Namely, with a subspace

$\mathfrak{N} \supset \text{ran}(T - T^*)$  define

$$\text{dom } A_0 = \mathfrak{H} := \mathcal{H} \ominus \mathfrak{N}, \quad A_0 = T \upharpoonright \text{dom } A_0.$$

Then  $\text{dom } A_0 \subset \ker(T - T^*)$ .

With  $A_0$  and  $K \in \mathbf{L}(\mathfrak{D}_A, \mathfrak{N})$  as in (7.1), the formula (4.1) provides a one-to one correspondence between all contractions  $X \in \mathbf{L}(D_{K^*})$  and all *qsc*-extensions of  $A$ . The operator form of all *qsc*-extensions with their resolvents was obtained in [7], [8]. Clearly, the subspaces  $\mathfrak{H}'$  and  $\mathfrak{H}'' = \mathfrak{H} \ominus \mathfrak{H}'$ , where

$$(7.2) \quad \mathfrak{H}' := \overline{\text{span}} \{ (T - zI)^{-1} \mathfrak{N} : |z| > 1 \} = \overline{\text{span}} \{ T^n \mathfrak{N} : n \in \mathbb{N}_0 \},$$

are invariant with respect to  $T$  and  $T^*$ , respectively. The inclusion  $\mathfrak{N} \subset \mathfrak{H}'$  implies that  $\mathfrak{H}'' \subset \mathfrak{N}^\perp = \text{dom } A \subset \ker(T - T^*)$ . Therefore the restriction of  $T^*$  to  $\mathfrak{H}''$  is a selfadjoint operator in  $\mathfrak{H}''$ . The restriction  $T \upharpoonright \mathfrak{H}' (= P_{\mathfrak{H}'} T \upharpoonright \mathfrak{H}')$  is called the  *$\mathfrak{N}$ -minimal part of  $T$* . Moreover,  $T$  is said to be  *$\mathfrak{N}$ -minimal* if the equality  $\mathfrak{H} = \mathfrak{H}'$  holds [4]. The subspaces  $\mathfrak{H}'$  and  $\mathfrak{H}^s$  of  $\mathcal{H} = \mathfrak{N} \oplus \mathfrak{H}_0$  defined in (7.2) and (4.5), respectively, are connected by  $\mathfrak{H}' = \mathfrak{N} \oplus \mathfrak{H}^s$ . Every *qsc*-extension  $T$  of the Hermitian contraction  $A_0$  in  $\mathcal{H}$  with  $\text{dom } A_0 = \mathfrak{H}$  generates a *pqs*-system in the following manner: let (1.7) be the block-operator representation of  $T$ , then the system  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  is a *pqs*-system.

**Proposition 7.1.** *Let  $A_0$  be a Hermitian contraction in  $\mathcal{H} = \mathfrak{N} \oplus \mathfrak{H}$  with  $\mathfrak{H} = \text{dom } A_0$ , let  $T$  be a *qsc*-extension of  $A$  in  $\mathcal{H}$ , and let  $\tau$  be the *pqs*-system generated by  $T$  with the state space  $\mathfrak{H}$  and the input and the output space  $\mathfrak{N}$ . Then the following statements are equivalent:*

- (i) *the Hermitian contraction  $A_0$  is simple;*
- (ii) *the *qsc*-extension  $T$  of  $A_0$  is  $\mathfrak{N}$ -minimal;*
- (iii) *the *pqs*-system  $\tau$  is minimal.*

*Proof.* The equivalence (i)  $\Leftrightarrow$  (ii) was proved in [4]. For the proof of (i)  $\Leftrightarrow$  (iii), observe that the simplicity of  $A_0$  is equivalent to  $\mathfrak{H}_0^s = \mathfrak{H}$ . Since  $\text{ran } K^* \subset \mathfrak{D}_{A_0}$  and  $\mathfrak{D}_{A_0}$  is invariant under  $A_0$ , the inclusion  $\mathfrak{H}_0^s \subset \mathfrak{D}_{A_0}$  holds. Hence,  $\mathfrak{H}_0^s = \mathfrak{H}$  implies in fact that  $\ker D_{A_0} = \{0\}$ , i.e.,  $\|Af\| < \|f\|$  holds for all  $f \in \mathfrak{H} \setminus \{0\}$ . Now the assertion follows from Proposition 4.2.  $\square$

**7.2. Transfer functions and  $Q$ -functions.** Let  $T$  be a *qsc*-operator in a separable Hilbert space  $\mathcal{H}$  and let  $\mathfrak{N}$  be a subspace of  $\mathfrak{H}$  such that  $\mathfrak{N} \supset \text{ran}(T - T^*)$ . The operator-valued function

$$(7.3) \quad Q_T(z) = P_{\mathfrak{N}}(T - zI)^{-1} \upharpoonright \mathfrak{N}, \quad |z| > 1,$$

is said to be a  *$Q$ -function* of  $T$ , cf. [4]. Analytical properties of the  *$Q$ -function* of *qsc*-operators and its applications to the parametrization of the resolvents of all *qsc*-extensions of corresponding Hermitian contraction were established in [4].

Let  $\mathfrak{N}$  be a Hilbert space. An operator-valued function  $Q(z)$  with values in  $\mathbf{L}(\mathfrak{N})$  and holomorphic outside  $\mathbb{D}$  is said to belong to the class  $\mathbf{Q}(\mathfrak{N})$  (see [4, Section 6]) if

(S1)  $Q(z)$  has the asymptotic expansion

$$(7.4) \quad Q(z) = -\frac{1}{z}I + \frac{1}{z^2}F + o\left(\frac{1}{z^2}\right), \quad z \rightarrow \infty;$$

(S2) the kernel

$$\frac{Q(z) - Q^*(\xi) - Q^*(\xi)(F - F^*)Q(z)}{z - \bar{\xi}}$$

is nonnegative;

(S3) the kernel

$$\frac{(1 - z^2)Q(z) - (1 - \bar{\xi}^2)Q^*(\xi) - (1 - z\bar{\xi})Q^*(\xi)(F - F^*)Q(z) - (z - \bar{\xi})I}{z - \bar{\xi}}$$

is nonnegative;

(S4) there exist a complex number  $z_0$ ,  $|z_0| > 1$ , and a vector  $f \in \mathfrak{N}$ , such that

$$\frac{Q(z_0) - Q^*(z_0) - Q^*(z_0)(F - F^*)Q(z_0)}{z_0 - \bar{z}_0} \neq Q^*(z_0)Q(z_0)f.$$

If  $T$  is a  $qsc$ -operator in the Hilbert space  $\mathfrak{H}$ ,  $\mathfrak{N}$  is a subspace of  $\mathfrak{H}$  such that  $\text{ran}(T - T^*) \subset \mathfrak{N}$ , and  $Q_T(z)$  is its  $Q$ -function defined by (7.3), then the function  $Q_T(z)$  belongs to the class  $\mathbf{Q}(\mathfrak{N})$ , see [4]. The converse statement is also true.

**Theorem 7.2** ([4]). *Let  $Q(z)$  belong to  $\mathbf{Q}(\mathfrak{N})$ . Then there exist Hilbert spaces  $\mathcal{H} \supset \mathfrak{N}$ ,  $\mathfrak{N} \neq \mathcal{H}$ , and a  $\mathfrak{N}$ -minimal  $qsc$ -operator  $T$  in  $\mathfrak{H}$ , such that  $\mathfrak{N} \supset \ker(T - T^*)$ , and*

$$Q(z) = P_{\mathfrak{N}}(T - zI)^{-1} \upharpoonright \mathfrak{N}, \quad |z| > 1.$$

The next proposition gives connections between the transfer function of a  $pqs$ -system  $\tau$  and the  $Q$ -function of the corresponding  $qsc$ -operator  $T$ .

**Proposition 7.3.** *Let  $\tau = \{T; \mathfrak{N}, \mathfrak{N}, \mathfrak{H}\}$  be a  $pqs$ -system. Then the transfer function  $\Theta(\lambda)$  of  $\tau$  and the  $Q$ -function of the  $qsc$ -operator  $T$  are connected by the following relations*

$$(7.5) \quad Q(z) = \left( \Theta \left( \frac{1}{z} \right) - zI_{\mathfrak{N}} \right)^{-1}, \quad |z| > 1; \quad \Theta(\lambda) = \frac{1}{\lambda}I + Q^{-1} \left( \frac{1}{\lambda} \right), \quad \lambda \in \mathbb{D}.$$

*Proof.* With  $W(z) = \Theta(1/z) - zI$ ,  $|z| > 1$ , the resolvent  $(T - zI)^{-1}$  has the form

$$\begin{pmatrix} W^{-1}(z) & -W^{-1}(z)C(A - zI)^{-1} \\ -(A - zI)^{-1}BW^{-1}(z) & (A - zI)^{-1}(I + BW^{-1}(z)C(A - zI)^{-1}) \end{pmatrix}.$$

Here the Schur-Frobenius formula has been applied; cf. [4, Section 2.4]. It follows that

$$P_{\mathfrak{N}}(T - zI)^{-1} \upharpoonright \mathfrak{N} = W^{-1}(z).$$

Thus, the relations (7.5) hold. □

Observe that Theorem 5.1 and Theorem 7.2 imply that  $\Theta(\lambda)$  belongs to  $\mathbf{S}^{qs}(\mathfrak{N})$  if and only if

$$Q(z) = \left( \Theta \left( \frac{1}{z} \right) - zI_{\mathfrak{N}} \right)^{-1}, \quad |z| > 1,$$

belongs to  $\mathbf{Q}(\mathfrak{N})$ .

**7.3. Scalar functions of the class  $\mathbf{S}^{qs}$  and Jacobi matrices.** Let  $l_2(\mathbb{N})$  and  $l_2(\mathbb{N}_0)$  be the Hilbert spaces of square summable complex-valued sequences

$$x = \{x_1, x_2, \dots, x_k, \dots\}, \quad x = \{x_0, x_1, \dots, x_k, \dots\},$$

considered as semi-infinite vector-columns, with the inner product given by

$$(x, y) = \sum_{k=1}^{\infty} x_k \bar{y}_k, \quad (x, y) = \sum_{k=0}^{\infty} x_k \bar{y}_k,$$

respectively. Clearly  $\mathbb{C} \oplus l_2(\mathbb{N}) = l_2(\mathbb{N}_0)$ . Define the vectors  $\{\delta_k\}$ ,  $k \in \mathbb{N}_0$ , by

$$\delta_0 = (1, 0, 0, \dots)^T, \quad \delta_k = (0, \dots, 0, 1, 0, 0, \dots)^T, \quad k \in \mathbb{N},$$

so that 1 is the  $(k+1)$ -st entry. Then the vectors  $\{\delta_k\}$  form an orthonormal basis in  $l_2(\mathbb{N}_0)$ .

**Theorem 7.4.** *Let the scalar function  $\Theta(\lambda)$  belong to the class  $\mathbf{S}^{qs}$ .*

- (i) *If  $\Theta(\lambda)$  is rational with  $n$  poles then any minimal pqs-system  $\tau = \{T, \mathbb{C}, \mathbb{C}, \mathfrak{H}\}$  with the transfer function  $\theta(\lambda)$  is unitarily equivalent to the pqs-system*

$$\tau_0 = \{T_0, \mathbb{C}, \mathbb{C}, \mathbb{C}^n\},$$

*where the operator  $T_0$  in the Hilbert space  $\mathbb{C} \oplus \mathbb{C}^n = \mathbb{C}^{n+1}$  with respect to the canonical basis  $\{\delta_k\}_{k=0}^n$  is given by the three-diagonal Jacobi matrix*

$$(7.6) \quad T_0 = \begin{pmatrix} \Theta(0) & a_0 & 0 & 0 & \cdot & \cdot & \cdot \\ a_0 & b_1 & a_1 & 0 & \cdot & \cdot & \cdot \\ 0 & a_1 & b_2 & a_2 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & a_{n-1} \\ \cdot & \cdot & \cdot & \cdot & 0 & a_{n-1} & b_n \end{pmatrix}.$$

- (ii) *If  $\Theta(\lambda)$  is not rational then any minimal pqs-system  $\tau = \{T, \mathbb{C}, \mathbb{C}, \mathfrak{H}\}$  with the transfer function  $\Theta(\lambda)$  is unitarily equivalent to the pqs-system*

$$\tau_0 = \{T_0, \mathbb{C}, \mathbb{C}, l_2(\mathbb{N})\},$$

*where the operator  $T_0$  in the Hilbert space  $\mathbb{C} \oplus l_2(\mathbb{N}) = l_2(\mathbb{N}_0)$  with respect to the canonical basis  $\{\delta_k\}_{k=0}^{\infty}$  is given by the semi-infinite three-diagonal Jacobi matrix*

$$(7.7) \quad T_0 = \begin{pmatrix} \Theta(0) & a_0 & 0 & 0 & 0 & \cdot & \cdot \\ a_0 & b_1 & a_1 & 0 & 0 & \cdot & \cdot \\ 0 & a_1 & b_2 & a_2 & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

*In both cases  $a_k > 0$  and  $b_k \in \mathbb{R}$  for all relevant  $k$ , and these numbers are uniquely determined by the function  $\Theta(\lambda)$ .*

*Proof.* Let  $\Theta(\lambda) \in \mathbf{S}^{qs}$ . By Proposition 7.3 the function  $Q(z) = (\Theta(1/z) - z)^{-1}$ ,  $|z| > 1$ , belongs to the class  $\mathbf{Q}$ . Without loss of generality it can be assumed that  $\text{Im } \Theta(0) \geq 0$ . Since the function  $w(\lambda) = \Theta(\lambda) - \Theta(0)$  belongs to the Herglotz-Nevanlinna class, the function  $Q(z)$  has a holomorphic continuation to the lower half-plane. If  $\Theta(\lambda)$  is rational

with  $n$  poles  $\{\mu_k\}$ , then these poles are simple and belong to  $(-\infty, -1) \cup (1, \infty)$ . It follows that

$$\Theta\left(\frac{1}{z}\right) = a \frac{\prod_{k=1}^n (z - \lambda_k)}{\prod_{k=1}^n (z - \mu_k)}, \quad \operatorname{Im} a \geq 0,$$

and  $Q(z)$  is rational with poles of total multiplicity  $n + 1$ , in the upper half-plane.

Let  $\tau = \{T, \mathbb{C}, \mathbb{C}, \mathfrak{H}\}$  be a minimal  $pqs$ -system with the transfer function  $\Theta(\lambda)$ . Then  $Q(z)$  is the  $Q$ -function of the corresponding  $qsc$ -operator  $T$ , i.e.,

$$Q(z) = ((T - zI)^{-1}\bar{1}, \bar{1}), \quad |z| > 1,$$

where

$$\bar{1} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix}, \quad 1 \in \mathbb{C}, \quad \mathbf{0} \in \mathfrak{H} \quad (\text{null-vector}).$$

The operator  $T$  in the Hilbert space  $\mathcal{H} = \mathbb{C} \oplus \mathfrak{H}$  is dissipative ( $\operatorname{Im} T(f, f) \geq 0$  for all  $f \in \mathcal{H}$ ),  $\dim \operatorname{ran} (T - T^*) \leq 1$ , and by Proposition 7.1 the operator  $T$  is  $\mathbb{C}$ -minimal, i.e.,  $\overline{\operatorname{span}} \{T^n \bar{1} : n \in \mathbb{N}_0\} = \mathcal{H}$ . If  $\operatorname{Im} \Theta(0) = 0$ , then  $T$  is selfadjoint with the cyclic vector  $\bar{1}$ . If  $\operatorname{Im} \Theta(0) \neq 0$ , then  $T$  is a prime dissipative operator with a rank-one imaginary part and, moreover,  $\operatorname{ran} (T - T^*) = \mathbb{C} \oplus \{0\}$ .

Note that  $T$  is unitarily equivalent to  $T_0$  given by (7.6) (in the case of a rational  $Q(z)$ ) or by (7.7) (in the opposite case) with respect the orthonormal basis  $\{\delta_k\}$ , i.e., there exists a unitary operator  $U \in \mathbf{L}(\mathcal{H}, \mathbb{C}^{n+1})$  or  $U \in \mathbf{L}(\mathcal{H}, l_2(\mathbb{N}_0))$  such that  $UT = T_0U$ ; cf. [9, Theorem 2.10, Theorem 6.1, Remark 6.2]. Moreover,

$$(7.8) \quad U\bar{1} = \delta_0.$$

Thus, for  $|z| > 1$  it follows that  $Q(z) = ((T_0 - zI)^{-1}\delta_0, \delta_0)$ . The entries of the matrix  $T_0$  can be found by the continued-fraction expansion of the function  $Q(z)$ :

$$Q(z) = \frac{-1}{z - \Theta(0)} + \frac{-a_0^2}{z - b_1} + \frac{-a_1^2}{z - b_2} + \dots + \frac{-a_{n-1}^2}{z - b_n} + \dots,$$

Note that  $T_0$  is selfadjoint with the cyclic vector  $\delta_0$  if  $\operatorname{Im} \Theta(0) = 0$ , and that  $T_0$  is a prime dissipative operator with a rank-one imaginary part and  $\operatorname{ran} (T_0 - T_0^*) = \operatorname{span} \{\delta_0\}$  if  $\operatorname{Im} \Theta(0) \neq 0$ . From (7.7) it follows that

$$T_0 = \begin{pmatrix} D_0 & B_0^* \\ B_0 & A_0 \end{pmatrix} : \begin{pmatrix} \mathbb{C} \\ \mathbb{C}^n \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ \mathbb{C}^n \end{pmatrix} \quad \text{or} \quad T_0 = \begin{pmatrix} D_0 & B_0^* \\ B_0 & A_0 \end{pmatrix} : \begin{pmatrix} \mathbb{C} \\ l_2(\mathbb{N}) \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ l_2(\mathbb{N}) \end{pmatrix},$$

respectively, where  $D_0 = \Theta(0)$ ,

$$B_0 1 = \begin{pmatrix} a_0 \\ 0 \\ 0 \\ \vdots \end{pmatrix}, \quad B_0^* \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \end{pmatrix} = a_0 x_1 \delta_0,$$

and

$$A_0 \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = \begin{pmatrix} b_1 & a_1 & 0 & 0 & \cdot & \cdot \\ a_1 & b_2 & a_2 & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix}.$$

Decompose  $T$  according to  $\mathcal{H} = \mathbb{C} \oplus \mathfrak{H}$ :

$$T = \begin{pmatrix} D & B^* \\ B & A \end{pmatrix} : \begin{pmatrix} \mathbb{C} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ \mathfrak{H} \end{pmatrix}.$$

Because (7.8) holds, the unitary operator  $U$  takes the following block operator matrix form

$$U = \begin{pmatrix} 1 & 0 \\ 0 & V \end{pmatrix} \begin{pmatrix} \mathbb{C} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ \mathbb{C}^n \end{pmatrix} \quad \text{or} \quad U = \begin{pmatrix} 1 & 0 \\ 0 & V \end{pmatrix} \begin{pmatrix} \mathbb{C} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ l_2(\mathbb{N}) \end{pmatrix},$$

respectively. Hence,  $VA = A_0V$ ,  $VB = B_0$ , i.e., the  $pqs$ -systems  $\tau$  and  $\tau_0$  are unitarily equivalent.  $\square$

Here is a simple example to illustrate the situation.

**Example 7.5.** Consider the scalar-valued function  $\Theta(\lambda)$  defined by

$$\Theta(\lambda) = d + \frac{1 - \sqrt{1 - \lambda^2}}{2\lambda}, \quad \lambda \in \text{Ext} \{(-\infty, -1] \cup [1, +\infty)\}.$$

Then  $\Theta(0) = d$  and  $W(\lambda) = \Theta(\lambda) - \Theta(0)$  is given by

$$W(\lambda) = \frac{1 - \sqrt{1 - \lambda^2}}{2\lambda}, \quad \lambda \in \text{Ext} \{(-\infty, -1] \cup [1, +\infty)\}.$$

Clearly  $W(\lambda)$  is a Herglotz-Nevanlinna function. It follows that

$$-2W\left(\frac{1}{z}\right) = \sqrt{z^2 - 1} - z = \frac{1}{\pi} \int_{-1}^1 \frac{\sqrt{1 - t^2} dt}{t - z}, \quad z \in \text{Ext} [-1, 1].$$

Hence  $W(1) = -W(-1) = 1/2$ . Moreover,  $\Theta(\lambda)$  belongs to the class  $\mathbf{S}^{qs}$  if and only if  $|d| \leq 1/2$ ; see (4.12). Assume that this condition is satisfied. Consider the weighted Hilbert space  $L_2([-1, 1], \rho(t))$  with the weight function

$$\rho(t) = \frac{2}{\pi} \sqrt{1 - t^2}, \quad t \in [-1, 1].$$

Define the operator  $A$  in  $L_2([-1, 1], \rho(t))$  by

$$(Af)(t) = tf(t), \quad f(t) \in L_2([-1, 1], \rho(t)).$$

Then  $A$  is a selfadjoint contraction. The function  $e_0(t) = 1$ ,  $t \in [-1, 1]$  belongs to  $L_2([-1, 1], \rho(t))$  and  $\|e_0\| = 1$ . Define the operator  $B : \mathbb{C} \rightarrow L_2([-1, 1], \rho(t))$  by

$$Bc = \frac{1}{2} c e_0(t), \quad c \in \mathbb{C}.$$

Then

$$B^*f(t) = \frac{1}{\pi} \int_{-1}^1 f(t) \sqrt{1 - t^2} dt, \quad f(t) \in L_2([-1, 1], \rho(t)).$$

Let  $D$  be the multiplication by  $d$  in the space  $\mathbb{C}$ . One can check that

$$T = \begin{pmatrix} D & B^* \\ B & A \end{pmatrix} : \begin{pmatrix} \mathbb{C} \\ L_2([-1, 1], \rho(t)) \end{pmatrix} \rightarrow \begin{pmatrix} \mathbb{C} \\ L_2([-1, 1], \rho(t)) \end{pmatrix}$$

is a *qsc*-operator. Moreover, the corresponding *pqs*-system

$$\tau = \{T; \mathbb{C}, \mathbb{C}, L_2([-1, 1], \rho(t))\}$$

is minimal and has the transfer function  $\Theta(\lambda)$ . Since the operator  $A$  is unitarily equivalent to the Jacobi matrix (see [18])

$$A_0 = \begin{pmatrix} 0 & 1/2 & 0 & 0 & \cdot & \cdot & \cdot \\ 1/2 & 0 & 1/2 & 0 & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix},$$

the unitarily equivalent three-diagonal minimal *pqs*-system is of the form

$$T_0 = \begin{pmatrix} d & 1/2 & 0 & 0 & 0 & \cdot & \cdot \\ 1/2 & 0 & 1/2 & 0 & 0 & \cdot & \cdot \\ 0 & 1/2 & 0 & 1/2 & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

## REFERENCES

- [1] Yu.M. Arlinskiĭ, Contractive extensions of a dual pair of contractions and their resolvents, *Ukrain. Math. J.*, 37 no. 2 (1985), 247–250.
- [2] Yu.M. Arlinskiĭ, A class of contractions in a Hilbert space, *Ukrain. Math. J.*, 39 no. 6 (1987), 691–696.
- [3] Yu.M. Arlinskiĭ, S. Hassi, and H.S.V. de Snoo,  $Q$ -functions of Hermitian contractions of Kreĭn–Ovcharenko type, *Integral Equations Operator Theory*, 53 no. 2 (2005), 153–189.
- [4] Yu.M. Arlinskiĭ, S. Hassi, and H.S.V. de Snoo,  $Q$ -functions of quasi-selfadjoint contractions, *Oper. Theory Adv. Appl.*, 163 (2006), 23–54.
- [5] Yu.M. Arlinskiĭ, S. Hassi, and H.S.V. de Snoo, Parametrization of contractive block operator matrices and passive discrete time systems, *Complex Analysis Operator Theory*, 1 (2007), 211–233.
- [6] Yu.M. Arlinskiĭ and E.R. Tsekanovskiĭ, Nonselfadjoint contractive extensions of Hermitian contractions and M.G. Kreĭn’s theorems, *Uspehi mat. nauk*, 37 (1982), 131–132 (Russian) [English transl.: *Russian Mathematical Surveys*, 37 (1982), 151–152].
- [7] Yu.M. Arlinskiĭ and E.R. Tsekanovskiĭ, Generalized resolvents of quasi-selfadjoint contracting extensions of a Hermitian contraction, *Ukrain. Mat. J.*, 35 (1983), no. 5 (Russian).
- [8] Yu.M. Arlinskiĭ and E.R. Tsekanovskiĭ, Quasi-selfadjoint contractive extensions of Hermitian contractions, *Teor. Funkts., Funkts. Anal. Prilozhen*, 50 (1988), 9–16 (Russian) [English transl.: *J. Soviet Math.*, 49 (1990), 1241–1247].
- [9] Yu. Arlinskiĭ and E.R. Tsekanovskiĭ, Non-self-adjoint Jacobi matrices with a rank-one imaginary part, *J. Functional Analysis*, 241 (2006), 383–438.
- [10] D.Z. Arov, Passive linear stationary dynamical systems. *Sibirsk. Math. J.*, 20 (1979), 211–228 (Russian) [English transl.: *Siberian Math. J.*, 20 (1979), 149–162].
- [11] D.Z. Arov, Stable dissipative linear stationary dynamical scattering systems. *J. Operator Theory*, 1 (1979), 95–126 (Russian) [English transl.: *Interpolation Theory, Systems, Theory and Related Topics. The Harry Dym Anniversary Volume*, *Oper. Theory Adv. Appl.*, 134 (2002), Birkhauser Verlag].
- [12] D.Z. Arov, M.A. Kaashoek, and D.P. Pik, Minimal and optimal linear discrete time-invariant dissipative scattering systems. *Integral Equations Operator Theory*, 29 (1997), 127–154.
- [13] D.Z. Arov, M.A. Kaashoek, and D.P. Pik, The Kalman–Yakubovich–Popov inequality for infinite dimensional discrete time dissipative system, *J. Operator Theory* (to appear).
- [14] D.Z. Arov and M.A. Nudel’man, A criterion for the unitary similarity of minimal passive systems of scattering with a given transfer function, *Ukrain. Math. J.*, 52 (2000), 161–172 (Russian).

- [15] D.Z. Arov and M.A. Nudel'man, Tests for the similarity of all minimal passive realizations of a fixed transfer function (scattering and resistance matrix), *Mat. Sb.*, 193 no. 6 (2002), 3–24 (Russian).
- [16] D.Z. Arov and O.J. Staffans, Bi-inner dilations and bi-stable passive scattering realizations of Schur class operator-valued functions (submitted).
- [17] M. Bakonyi and T. Constantinescu, *Schur's algorithm and several applications*, Pitman Research Notes in Mathematics Series, Longman House, Harlow, UK, 1992.
- [18] Yu. M. Berezanskii, *Expansions in Eigenfunctions of Self-Adjoint Operators*, Transl. Math. Monographs. 17, Amer. Math. Soc., Providence, RI, 1968.
- [19] S.S. Boiko and V.K. Dubovoj, On some extremal problem connected with the suboperator of scattering through inner channels of the system, Report of National Academy of Sciences of Ukraine, no. 4 (1997), 7–11.
- [20] S.S. Boiko, V.K. Dubovoj, B. Fritzsche, and B. Kirstein, Contraction operators, defect functions and scattering theory, *Ukrain. Math. J.*, 49 no. 4 (1997), 529–538.
- [21] S.S. Boiko, V.K. Dubovoj, B. Fritzsche, and B. Kirstein, Models of contractions constructed from the defect function of their characteristic function, *Oper. Theory Adv. Appl.*, 123 (2001), 67–87.
- [22] V.K. Dubovoj, R.K. Mohhamed, Defect functions of holomorphic contractive matrix functions, regular extensions and open systems, *Math. Nachr.*, 160 (1993), 69–110.
- [23] B. Fritzsche, B. Kirstein, and J. Lorenz, On an extension problem for contractive block Hankel operator matrices. *Analysis*, 25 (2005), 23–54.
- [24] M.G. Krein and I.E. Ovcharenko, On Q-functions and sc-extensions of a Hermitian contraction with nondense domain, *Sibirsk. Math. J.*, 18 (1977) 18, 1032–1056 (Russian) [English transl.: *Siberian Math. J.*, 18 (1977), 728–746].
- [25] F. Riesz and B. Sz.-Nagy, *Leçons d'Analyse Fonctionnelle*, Gauthier-Villars, 1965.
- [26] W. Rudin, *Real and complex analysis*, McGraw-Hill Book Co., New York, 1987.
- [27] B. Sz.-Nagy and C. Foias, *Harmonic analysis of operators on Hilbert space*, North-Holland, New York, 1970.

DEPARTMENT OF MATHEMATICAL ANALYSIS,, EAST UKRAINIAN NATIONAL UNIVERSITY,, KVAR-TAL MOLODYIOZHNY 20-A,, LUGANSK 91034,, UKRAINE

*E-mail address:* yma@snu.edu.ua

DEPARTMENT OF MATHEMATICS AND STATISTICS,, UNIVERSITY OF VAASA,, P.O. Box 700,, 65101 VAASA,, FINLAND

*E-mail address:* sha@uwasa.fi

DEPARTMENT OF MATHEMATICS AND COMPUTING SCIENCE,, UNIVERSITY OF GRONINGEN,, P.O. Box 800,, 9700 AV GRONINGEN,, NEDERLAND

*E-mail address:* desnoot@math.rug.nl